



Journal of Agriculture Sustainability and Environment

ISSN: 2997-271X

Global Scientific Research

www.jescae.com

Journal of Agriculture Sustainability and Environment

Vol.2, No.1

June, 2023

Chief Editor	Dr. Abdullah Khan
Edited by	Global Scientific Research
Published by	Global Scientific Research
Address:	GLOBAL SCIENTIFIC RESEARCH (LIMITED COMPANY): 8206 Louisiana Blvd Ne, Ste A #2531 Albuquerque, New Mexico 87113, United States
Journal Link:	https://www.jescae.com/index.php/jase
Email	journals.gsr@gmail.com
Website	www.jescae.com

CONTENTS

S.NO	TITLE	AUTHORS	PAGES
1	Conservation Tillage: A Sustainable Approach for Carbon Sequestration and Soil Preservation. A Review	Waqar Ahmad Khan, Gang Wang	1-24
2	Assessments on the Impacts of Climate Change on Food Production, Nutrition, Quality, and Resource Use Efficiency: A Review	Alemu Andualem, Tamirat Wato, Tariku Goa, Natnale Sitotaw, Gutema Urgi	25-34
3	Urinary outputs of nickel in association with their concentration levels in water, soil, and selected foods among farmers in the industrial estate of district swabi	Ijaz Ahmad, Niamat Ullah, Zia-ud-Din	35-48
4	A review of agroforestry as a sustainable and resilient agriculture	Asif Raihan	49-72

REVIEW ARTICLE

Conservation Tillage: A Sustainable Approach for Carbon Sequestration and Soil Preservation. A Review

Khan Waqar Ahmad ^{1*}, Gang Wang¹

¹School of Biological and Agricultural Engineering, Jilin University, Changchun 130022, China

Corresponding Author: Khan Waqar Ahmad: waqarahmadkhan1990@gmail.com

Received: 23 February, 2023 Accepted: 27 March 2023, Published: 29 March, 2023

Abstract

Minimum tillage is a soil conservation tillage aimed at minimizing soil disturbance required for productive crop production. Unlike intense tillage, which uses ploughs to alter the soil's structure, this tillage technique does not turn the soil over. Only secondary tillage is used sparingly in minimum tillage, with primary tillage being totally avoided. Practices like minimum furrowing, using organic fertilizer, using biological pest control techniques, and using less pesticides are all included in minimum tillage. Soil erosion and soil degradation have been increased by the use of conventional agricultural techniques, such as extensive tillage centered on the removal of crop residue. Global interest in finding various sustainable ways to lower the concentration of greenhouse gases in the atmosphere has grown in recent years as a result of the gradual increase in their concentration. The amount of carbon stored in soil is 2-4 times greater than that in the atmosphere and four times greater than that in vegetation. In order to prevent or, carbon sequestration (CS) delay dangerous climate change entails storing other forms of carbon or transferring CO₂ from the atmosphere into the soil. The potential of soils to store carbon and reduce the accelerated greenhouse effects by implementing various agricultural management strategies is covered in the current review. Conservation tillage techniques improve carbon sequestration in agricultural soils. Conservation tillage can be a practical way to store carbon in the soil and minimize the effects of climate change. Conservation tillage reduced the green house gas. Zero tillage has been identified as the most environmentally friendly tillage practice for the mitigation and adaptation to climate change processes. No-till farming is thought to make it possible to increase crop production sustainably in order to fulfill future agricultural demands.

Keywords: climate change; carbon sequestration; soil conservation; global warming; zero tillage; Soils degradation

Introduction

The growing concern for food security through improved soil management techniques demands identification of an environmentally friendly and crop yield sustainable system of tillage. Tillage is defined as the mechanical manipulation of the soil for the purpose of crop production affecting significantly the soil characteristics such as soil water conservation, soil temperature, infiltration and evapotranspiration processes. This suggests that tillage exerts impact on the soil purposely to produce crop and consequently affects the environment. As world population is increasing so the demand for food is increasing and as such the need to open more lands for crop production arises.

The yearning for yield increases to meet growing demand must be done in a way that soil degradation is minimal and the soil is prepared to serve as a sink rather than a source of atmospheric pollutants. Thus, conservation tillage, along with some complimentary practices such as soil cover and crop diversity (Corsi, Friedrich, Kassam, Pisante, & de Moraes Sà, 2012) has emerged as a viable option to ensure sustainable food production and maintain environmental integrity. This implies that conservation tillage is a component of conservation agriculture (CA).

Conservation Agriculture

define as a method of managing agro-ecosystems for improved and sustained productivity, increased profits and food security while preserving and enhancing the resource base and the environment. They added that minimum mechanical soil disturbance, permanent organic soil cover and crop diversification are the three basic principles of CA (Corsi *et al.*,2012). According to CTIC (2004), conservation tillage is any tillage system that leaves at least 30% of the soil surface covered with crop residue after planting to reduce soil erosion by water. (Lal *et al.*,1990) described conservation tillage as the method of seedbed preparation that includes the presence of residue mulch and an increase in surface roughness as the key criteria. Conservation tillage is an ecological approach to soil surface management and seedbed preparation. Conversion from conventional to conservation tillage, when this is done in line with the principle of CA, may improve soil structure, increase soil organic carbon, minimize soil erosion risks, conserve soil water, decrease fluctuations in soil temperature and enhance soil quality and its environmental regulatory capacity. Crop residue is an important and a renewable resource. Developing techniques for effective utilization of this vast resource is a major challenge. Improper uses of crop residues (e.g. removal, burning or ploughing under) can aid accelerated erosion, soil fertility depletion and environmental pollution through burning. The principle of conservation tillage involves maintenance of surface soil cover through retention of crop residues achievable by practicing zero tillage and minimal mechanical soil disturbance. Retention of crop residue protects the soil from direct impact of raindrops and sunlight while the minimal soil disturbance enhances soil biological activities as well as soil air and water movement. The aim of this review, therefore, was to examine the effects of conservation tillage on soil, crop and the net effect on the environment. This may provide farmers and other land users the information on the desirability of a conservation tillage system for sustainable crop yield increases with minimal negative impact on the soil and the environment.

Types of conservation tillage

Conservation tillage practices range from zero tillage (No-till), reduced (minimum) tillage, mulch tillage, ridge tillage to contour tillage. No tillage (NT) involves land cultivation with little or no soil surface disturbance, the only disturbance being during planting while minimum tillage means reduced level of soil manipulation involving ploughing using primary tillage implements. In mulch tillage, the soil is prepared or tilled in such a way that the plant residues or other materials are left to cover the surface to a maximum extent. Ridge tillage involves planting crops in rows either along both sides or on top of the ridges which are prepared at the commencement of the cropping season. When tillage is at right angles to the direction of the slope it is referred to as contour tillage. Table 1. Differences between conservation tillage and conventional tillage (Shahane et al., 2021)

Table 1. Differences between conservation tillage and conventional tillage (Shahane et al., 2021)

S.No.	Particulars	Conventional Tillage	Conservation Tillage
1	Soil health	Poor/degraded	Healthy soil
2	Tillage System	High Intensity plough based tillage system	Minimal tillage or zero tillage
3	Energy requirement	higher	Lower
4	Fallowing System	Ideal fallow land without any crop cover on soil surface	Growing of cover crops
5	sustainability	lower	Higher
6	Residue Management	Complete removal or burning of crop residue	Maintaining 30% soil surface covered with residues
7	Foot print on natural resources	Higher	Lower
8	Nutrient Management	Chemical based nutrient management or intensive use of chemical fertilizers	Integrated nutrient management with inclusion of organic sources and microbial inoculations
9	Fallowing System	Ideal fallow land without any crop cover on soil surface	Growing of cover crops

Conservation tillage and soil properties

Tillage impact is noticeable on soil physical, chemical and biological properties though in different magnitudes. Tillage impact also includes the effect on the soil environment in the form of runoff and soil erosion (Bhatt & Khera, 2006).

Soil physical properties

The effects of conservation tillage on soil properties are variable and depend on the specific system implemented. No-till (NT) systems, which maintain high soil coverage, have shown significant changes in soil properties, particularly in the upper few centimeters (Anikwe & Ubochi, 2007). Lal (1997a) suggests that soil physical properties generally favor no-till systems over traditional tillage-based systems. Many researchers have observed that NT improves both saturated and unsaturated hydraulic conductivity due to the continuity of pores (Benjamin, 1993) or flow through larger pores (Allmaras, Rickman, Ekin, & Kimball, 1977). Well-drained soils with lighter to medium textures and low humus content are particularly responsive to conservation tillage, especially no-till (Butorac, 1994).

NT technologies, according to Lal, Reicosky, and Hanson (2007), effectively reduce soil and crop residue disturbance, moderate soil evaporation, and minimize erosion losses. No-till soils also tend to exhibit more stable aggregates in the upper surface, resulting in higher total porosity. In a long-term experiment in Gottingen, Germany, Jacobs, Rauber, and Ludwig (2009) found that minimum tillage (MT) increased aggregate stability and concentrations of soil organic carbon (SOC) and nitrogen (N) in the upper 5-8 cm depth after several decades of tillage treatments.

In terms of water conservation, NT has shown greater effectiveness in humid and sub-humid tropics. Kargas, Kerkides, and Poulouvassilis (2012) found that untilled plots retain more water compared to tilled plots. Minimum tillage has been reported to improve the soil pore system, increasing storage pores and elongated transmission pores (Pagliai, Vignozzi, & Pellegrini, 2004). Higher water holding capacity and moisture content have also been

observed in the topsoil under NT compared to conventional ploughing (McVay et al., 2006). Therefore, replacing traditional tillage with conservation tillage has been proposed by many researchers to improve soil water storage and increase water use efficiency (WUE) (Fabrizzi et al., 2005, Silburn et al., 2007). Reduced tillage systems, including NT, have been found to result in greater water use efficiency compared to traditional tillage (McVay et al., 2006; Li, Huang, & Zhang, 2005). A study by Su et al. (2007) demonstrated that soil water storage and WUE were significantly higher in zero-tillage (ZT) than in conventional tillage (CT) over a six-year period.

In a study conducted in southwestern Nigeria, Busari and Salako (2012) observed higher unsaturated water flow parameters and infiltration rates under CT and MT than under ZT during the first year, but ZT showed higher infiltration parameters compared to CT by the end of the second year. This is because CT initially created fast-draining macro-pores (FDP) that facilitated infiltration, but these FDP decreased over time due to soil aggregate repackaging (Martínez, Fuentes, Silva, Valle, & Acevedo, 2008), resulting in lower infiltration rates under CT in the long term. Other studies (Pikul and Aase, 1995; Shukla et al., 2003) have also found higher infiltration rates under NT due to the protective effect of surface residue and the influence of SOC.

Additionally, less intense tillage practices not only preserve crop residue at the soil surface but also increase the activity of surface-feeding earthworms, creating numerous surface-connected macro-pores and inter-pedal voids, leading to higher rates of infiltration (Kemper, Trout, Segeren, & Bullock, 1987).

Aims and Objectives

1. To investigate the conservation agriculture is improve plant growth and soil health.
2. To discusses the potential of soils in sequestering carbon and mitigating the accelerated greenhouse effects by adopting different agricultural management practices.
3. To Investigate soils degradation soil degradation cause

Major Causes and Factors Affecting Soil Organic Carbon Depletion

The soil organic carbon pool is being quickly depleted as natural habitats give way to farmed ones. The amount of soil organic carbon pool depletion is 25–50% over 20–50 years in temperate temperature zones and 50–75% over 5–20 years in tropical climate zones after deforestation (Lal et al., 2004). When the C inputs in managed ecosystems (via crop residue retention, combined with the application of other biosolids) are greater than the outputs, the degree of depletion is minimal. The latter includes soil organic carbon losses due to leaching, mineralization, and increased erosion (caused by humans removing natural vegetation). By moving organic carbon-rich sediment from an agricultural land unit and surface runoff, agricultural soil erosion has been shown to disturb the global carbon cycle (Olson et al., 2016; Lal et al., 2009). Additionally, it has been noted that the emission of CO₂ and the depletion of the source of soil organic carbon are positively correlated (Navarro-Pedreño et al., 2021). According to reports, the loss of the soil organic carbon pool has a negative effect on soil quality and the equilibrium of nutrients and elements. Through runoff losses and high evaporation rates, it also affects the equilibrium of soil water and can cause a significant decline in soil biodiversity, including the activity of soil microbes. (Lal et al., 2004). asserts that deteriorating soil quality has a detrimental impact on net primary productivity and reduces the quantity and quality of plant biomass produced, leading to a significant depletion of the soil organic carbon pool.

Management of Soil Organic Carbon

The process of moving atmospheric carbon dioxide into the soil's C pool via humifying agricultural waste and other organic soil components (such biosolids), which are not quickly released back into the atmosphere, is known as soil

organic carbon sequestration (Olson et al., 2014). The main determinants of soil organic carbon sequestration are an increase in soil organic carbon content, its management through soil-based and crop-based management, applied by the use of C-enriched material (including mulches and biochar) prudent use of land resources and organic fertilizers. Low carbon agriculture is referred to as a sustainable method for reducing the effects of global warming, enhancing crop yields, and conserving the environment. Low-C agriculture practices are characterized by low GHG (including carbon dioxide) emissions and high soil organic carbon and vegetation storage. Utilizing best management practices to safeguard the environment, natural resources, and eventually crop productivity is the strategy's main objective. It is one of the best methods for reducing GHG emissions Zhang et al., 2017; Lal et al., 2018; De Gouvello et al., 2010; Gebara et al., 2013; de Magalhães et al., 2014). Table 2. Function of Soil organic Carbon (Shahane *et al.*, 2021)

Table 2. Function of Soil organic Carbon (Shahane *et al.*, 2021)

Plant	Improvement	Soil Maintenance	Reduce	Ecosystem
Increase in duration of shifting cultivation area available for cultivation	Enhanced the decomposition of soil pollutants Microbial population and diversity Biogeochemical cycling of nutrients	N/A	N/A	N/A
Crop Yield Improvement	Agregate Stability	Temperature	Bulk Density	Increase in Carbon Sequestration
Sustainability in Production System	Cation Exhange Capacity and base saturation	PH	Soil crusting and compaction	N/A
Quality improvement	Porosity	Soil Consistence	Erodibility and Erosion	Reduce greenhouse gas emission
Enhance resource and use efficiency	Infiltration	Air Circulation	Accumulation of toxic Material	Prevent station of tanks and enhance and their storage capacity and life
Profitability enhancement	Chelation of Micronutrients	Optium soil moisture	Reduce the leaching loss of nutrients	N/A
Reduced bioaccumulation of soil pollutants in the plants products	Water and nutrient retention capacity	Desirable soil structure spheriodal granular and crumby structure	N/A	N/A

Mechanism of Soil C Sequestration

The three main processes that lead to soil carbon sequestration are the formation of soil microaggregates, the soil's long-term stability, and the enhancement of soil structure through the deep insertion of SOC in the subsoil layers (Lal et al., 1997; Tisdall et al., 1982; Six, Bossuyt et al., 2000). The stability of macro-aggregates might protect soil organic matter (OM) from microbial activity. The concentration of clay and mineralogy have a major impact on aggregation. Furthermore, a positive relationship between aggregate size and total soil organic carbon content was shown by Beare *et al.*, 1994; Puget *et al.*, 2005). The way biomass C humifies is also influenced by soil properties, tillage practices, climate, and soil nutrient availability. The humification efficiency of biomass C is lower in warm, dry regions compared to cold, humid conditions. In addition, large surface area clayey soils perform better in terms of humification efficiency than coarse-textured soils. The no-till farming method had a positive effect on the effectiveness of humification. (Puget *et al.*, 2005) found that 8.3% of the total carbon in crop residue for plow-tillage crops and 11.9% for no-till practices was converted to soil organic carbon in maize crops grown in Coshocton, Ohio. In a separate study, (Allmaras *et al.*, 2004) discovered that humification was 26% more successful for no-till soils than it was for traditional tillage methods like using moldboard plows and chisels, which were reported to be 11% more effective. The availability of soil components such as N, P, S, Zn, and Cu affects the efficiency of humification because C is the main component of humus. (Himes *et al.*, 2018). found that 28 mg of carbon in 62 mg of oven-dry residue is needed to store the 10 mg of carbon in crop residue into 17.241 mg of humus. It also requires 143 kg S, 200 kg P, and 833 kg N, according to the writers. Consequently, for the leftover C to be humified, essential nutrients like Nitrogen, Phosphorous, and Sulfur must be present. Regarding this, (Jacinthe *et al.*, 2002).found that residue-C conversion into soil organic carbon for Luvisol in central Ohio was 32% when fertilizer treatment was applied, compared to 14% when it wasn't. Under the mulched soils, comparable soil organic carbon stocks (25.6 Mg C ha⁻¹) have been identified, both with and without fertilizer treatment. Nevertheless, additional SOC deposition only occurs in regions where more fertilizer was applied when mulching material is employed. The no-till approach does not greatly increase the soil organic carbon pool in the absence of adequate fertilizer (Campbell *et al.*, 2001). The amount of SOC sequestered is significantly influenced by the rates and locations of N fertilizer application (Gregorich *et al.*, 1995; Wanniarachchi *et al.*, 1999; Murungu *et al.*, 2011). The illuviation and translocation of C into the subsurface layers is another important mechanism. The results of the bioturbation generated by earthworms, termites, and the deep root system are climate changes and the translocation of deep C away from the anthropogenic zone (Lavelle *et al.*, 1989; Lorenz *et al.*, 2005).




Figure 1. Sequestering Carbon in Soils to Reduce Climate Change

Conventional Tillage and Soil Carbon Stocks (CS)

Developing a soil environment that is favourable to improved plant growth and development is the primary goal of any tillage technique. One of the key elements determining soil C stocks is tillage. SOM is significantly reduced as a result of the aggressive tillage techniques. While ground cover removal exposes the organic-rich topsoil layers to wind and water erosion, tillage exposes soils to air, which promotes SOM decomposition by soil bacteria (Wani et al., 2016). Moreover, soil microbial activity and the holes left by plant roots are disturbed by intensive tillage techniques. The SOM is rapidly degraded and lost as a result of the fast-mechanical cultivation; the SOM gets protected within the soil aggregates. Tillage techniques also cause the soil aggregates to break down, increasing the amount of oxygen available and the surface area that organic material is exposed to.

Physiochemical Properties of Conventional Tillage Practices

Poor Infiltration and excess runoff. Deep layers compaction and structure disability. Nutrient loss and reduced CEC. Salinization and acidification.

Biological Properties of Conventional Tillage Practices

Reduced diversity of soil organisms. Reduced enzymatic activity and affect nutrient cycling. Reduced number of plants associated microorganisms. (Hussain et al., 2021)

Conservation Agriculture and NT for SOC

A different approach to increasing agricultural output in a sustainable way has been mentioned: the conservation agriculture (CA) system. In agricultural environments, this technique is widely thought to increase infiltration rates, lessen erosion problems and improve soil quality and organic C levels (Kahlon et al., 2013). According to a different study by (Prasad et al., 2016). conservation agriculture lessens the problems with soil degradation associated with rainfed agriculture. Conservation agriculture includes crop mulching, proper crop rotation, and no-till farming, which involves less soil disturbance. (Somasundaram et al., 2020). Conservation agriculture, in contrast to traditional agriculture operations, primarily aims to maximize yields at the expense of the environment. (Dumanski et al., 2006) state that conservation agriculture involves the supply of modern agricultural technology to improve crop production and maintain the health and integrity of the eco-system. The FAO recognized that the CA system lessens the negative effects of climate change, improves sustainable land management, and improves crop productivity without endangering the environment. (Pisante et al., 2015; Pisante et al., 2012). Over the past few years, Californian agriculture (CA) has been more well-known because of its many advantages, which include improved soil fertility and water retention, long-term sustainable productivity, and the reduction of climate change (González-Sánchez et al., 2012; Palm et al., 2014; Busari et al., 2015). Contrary to traditional systems, conservation agricultural methods in highland crop production systems enhance soil water and nutrient status, residual water content, soil infiltration rate, and organic carbon content (Thierfelder et al., 2009; Ella et al., 2016). Three fundamental ideas underlying CA: minimizing soil disturbance through no-till practices, keeping soil cover with mulching, and modifying crop rotation and intercropping techniques. Using woody crops to increase yields in low-fertility soils without impacting the environment was also suggested by several researchers (Assessment et al., 2015). Based on integrated nutrient management, (Lal et al., 1990).

Conservation Tillage Enhanced the Biodiversity

In addition to reducing soil and nutrient erosion, conservation tillage techniques like RT and ZT improve soil microflora and faunal variety, SOC, and related soil characteristics (Das et al., 2019; Raj et al., 2022).

Soil Degradation

When a land-use system's potential productivity becomes detrimental and the land is unable to fulfill its environmental regulatory roles of absorbing, storing, and recycling nutrients, water, and energy, this is referred to as land degradation (Oldeman et al., 1992). Once more, the measured loss or decline of a soil's present or showed ability to yield plant materials of the right amount and quality is commonly referred to as soil degradation. Some scholars Blaikie et al., 2015; Chisholm et al., 1987; Blum et al., 2004). Contend that the phrase "land degradation" is more inclusive than "soil degradation." However, as land and soil are alike in the majority of soil management literature, the terms land degradation and soil degradation will be used synonymously in the parts that follow. Soil degradation is caused by a number of chemical, physical, and biological processes Lal et al., 2020; Eswaran et al., 2001). Crusting, hard setting, Deterioration of soil structure desertification, erosion, are some of the physical processes. fertility loss, Leaching, salinization, acidification and pollution are some of the chemical processes. The decrease in soil biodiversity and the depletion of carbon are two examples of the biological processes causing soil deterioration. A difference between land quality and land usage causes land deterioration, claims (Beinroth et al., 1994). In certain root-restrictive shallow soils in West Africa, yield decreases of 30–90% due to erosion have been reported by (Mbagwu et al., 1984; Lal et al., 1987). In Ohio and other Midwestern USA states, erosion decreased row crop yields by 20–40% (Fahnestock et al., 1996). In the Colombian Andes, (Schumacher et al., 1994) have noted significant losses on certain sites as a result of rapid erosion. Soil erosion and desertification have caused a 50% decrease in the production of some African farms (Dregne, 1990). Due to historical soil erosion, Africa's yield decline can vary from 2 to 40%, with a mean loss of 8.2% across the continent (Ruppenthal et al., 1995). Furthermore, Asia, which includes China, India, Israel, Iran, Lebanon, Jordan and Pakistan, Nepal, has significant productivity losses (20%) as a result of erosion (Lal et al., 1995). Over a seven-year period, agricultural product decreases in 20% for soybeans, Ohio are 25% for maize and 30% for oats. (Lal et al., 1996). Table 3. Impacts of soil deterioration on crop output and growth. (Shahane et al., 2021). Table 4. Characteristics of healthy soil (Shahane et al., 2021)

Table 3. Impacts of soil deterioration on crop output and growth. (Shahane et al., 2021)

S.No.	Crop	Soil degradation related problem	effect	Correction measure suggested
1	Wheat	Salinity due to irrigation water	decrease in wheat growth parameters, harvest index, and grain and straw yields as irrigation water's electric conductivity rises from 0.7 to 12 dS m ⁻¹	When Azospirillum sp. isolated from saline soil is used, wheat grain yield significantly increases over control.
2	Rice bean	Soil acidity	Soil acidity reduced crop growth and yield as well as	Utilizing lime at a rate of 0.6 t ha ⁻¹ improves all growth and yield

			economic metrics (gross and net return, B:C ratio, production efficiency, and economic efficiency).	characteristics, leading to an increase in yield of 0.42 t ha ⁻¹ , or 221.31 and 164.34 US dollars in gross and net returns ha ⁻¹ , respectively.
3	Rice	Acidity of soil (acid sulfate soil) and aluminum toxicity	decrease in rice output brought on by more aluminum toxicity and acidity in the soil; decreased availability of exchangeable cations (Ca, Mg, and K)	Positive effect of addition of amendments such as magnesium limestone, sugarcane based organic fertilizers and fused magnesium phosphate
4	Chickpea	Sensitivity of sodium salt (sodium chloride)	The rise in sodium chloride concentration has a considerable impact on vegetative and reproductive growth, or the quantity of flower buds and pods; the crop's podding stage was shown to be the most vulnerable.	N/A
5	Garden Pea	Acidic soil	Acidity of the soil has a negative impact on garden pea development and soil characteristics	Application of corn or lantana camera biochar (@ 6 to 18 t ha ⁻¹) had a positive impact on crop growth metrics. Improvements in the soil's total nitrogen, accessible

6	Pea	Acidity of soil	N/A	phosphorus, potassium concentration, and porosity following crop harvest. Lime application at 7.5 t ha ⁻¹ increased grain yield and dry matter production by 0.50–0.55 t ha ⁻¹ and 1.37–1.72 t ha ⁻¹ , respectively.
7	Wheat	Waterlogging	After 21 days of sowing, waterlogging for 15 days lowers wheat yields in neutral soil (pH of 7.0), salty soil (pH of 8.2), acidic soil (pH of 9.0), and sodic soil (pH of 9.4).	N/A
8	Rice	Saline sodic soil	Saline sodic soil's detrimental effects on plant growth and yield	Rice growth and yield parameters were significantly improved by the use of gypsum at 9.5 t ha ⁻¹ and irrigation spaced four days apart. There was also a considerable increase in rice grain and straw yield.
9	French Bean	Chemical degradation (nutrient deficiency) in acidic soil	reduced growth and yield characteristics as a result of infertile soil	Growth and yield attributes improved as a result of applying three primary nutrients at the

appropriate rate in
addition to boron.

Table 4. Characteristics of healthy soil(Shahane et al., 2021)

S. No	Attributes	Description
1	Resilience	Healthy soils can bounce back fast from adverse events like compaction.
2	Important function of healthy soil	Carbon cycles, nutrient cycles, preservation of soil structure, control of pests and diseases
3	Resistance to being degraded	Good tilth, internal drainage, low plant parasite populations, and these characteristics help soils resist the damaging impacts of compaction and wet durations.
4	Sufficient supply of nutrients although	For plants to flourish, there must be a sufficient supply of nutrients; at the end of the growing season, there shouldn't be an excessive amount of phosphorous and nitrogen left in highly soluble forms or enriching the soil's surface. The most likely times for fertilizer leaching and runoff are after crops are harvested and before the next crops are well-established.
5	Good soil tilth	Compared to soil with poor tilth, excellent tilth soil is less compacted, spongier, and allows roots to grow more fully. Water infiltration and storage for plant use later on is additionally supported by a soil with a stable and beneficial soil structure.
6	No chemicals that harm plants	Naturally occurring hazardous substances can include excess salts in arid areas or soluble aluminum in acidic soils. Human activity can introduce potentially dangerous chemicals through the application of sewage sludge containing high concentrations of toxic components or fuel-oil spills.
7	Low weed pressure	It is crucial to have minimal weeds so that the crop has less competition for nutrients, water, and light.
8	Sufficient depth	Full root system growth is supported by soils that are deep enough to contain a layer that can impede drainage and/or root development.
9	Good internal drainage	Soils that dry up rapidly can benefit from timely field operations. Additionally, for the best possible root health, oxygen needs to be able to enter the root zone, and proper drainage makes this possible.
10	high numbers of microbes that promote plant growth	Earthworms and a variety of bacteria, fungus, and actinomycetes are examples of organisms that aid in the cycling of nutrients and make them available to plants. Also, soil organisms generate compounds that stimulate plant growth.

Causes of Soil Degradation

Both natural and man-made factors can lead to soil degradation. Natural reasons include topographic and climatic elements like, frequent floods and tornadoes, steep slopes ,storms and strong winds, leaching in humid areas, intense rains and drought in arid areas. Anthropogenic causes of soil degradation include overextraction of ground water, shifting farming, desurfacing of the soil, excessive grazing, indiscriminate use of agrochemicals, and deforestation and overexploitation of vegetation.

Types of Soil Degradation

In 1991, ISRIC, in collaboration with FAO and UNEP, released a global map showing the state of soil degradation caused by human activity. A generic classification known as the GLASOD classification was created in advance of the map., Water erosion chemical deterioration, wind erosion, physical deterioration, and loss of biological activity are the five primary kinds of soil degradation, according to (Global Assessment of Soil Deterioration) (Oldeman *et al.*, 1992). Every type has multiple subtypes, with the exception of biological deterioration. The following lists these varieties and subtypes. Table 5. Soil degradation types and subtypes.




Figure 2. Land area so far degraded by different processes (Data) (Lal *et al.*, 1996)

Table 5. Soil degradation types and subtypes

Type	Subtypes
Water Erosion	Loss topsoil Terrain deformation/mass movement site effect Reservoir sedimentation Flooding Sea weed destruction
Wind Erosion	Loss topsoil Terrain deformation Overblowing
Chemical deterioration	Loss of Nutrient and organic matter Salination Acidification

	Eutrofication
Physical deterioration	Compaction, sealing, and crusting Water Logging Lowering of water table Subsidence of organic soils

Extent of Soil Degradation

Approximately 38% of the planet's agricultural land have been deemed degraded. Africa has 65% of degraded territory, Central America has 74%, and South America has 45%. There is a significantly lower percentage of damaged grassland and forests—21% and 18%, respectively. If we simply take into account land that has been utilized (forests, permanent pasture, and agricultural areas), the percentage of degraded land is 14% and the percentage of severely degraded land is 23%. The area impacted by human-induced soil degradation was judged to be mildly deteriorated in 38% of the cases (749 M ha), moderately degraded in 46% of cases (910 M ha), strongly degraded in 15% of cases (296 M ha), and extremely degraded in fewer than 1% of cases (9.3 M ha) (Lal et al., 1996) In Asia, nutrient imbalances in the soil, overfertilization, pollution, and soil loss processes have a negative impact on soil health and quality.

The soil's organic stuff is deteriorating daily. With the growth in the nation's population over the past few decades, it has grown increasingly intense. On Earth, 25% of all species are found in soil. Table 6. Indicators of soil health and their measurements (Shahane et al., 2021)

Table 6. Indicators of soil health and their measurements (Shahane et al., 2021)

Soil health indicator	Unit of measurement	Ideal values for health soil indicators (agricultural soil) and Method
Texture	twelve classes based on the relative proportion of sand, clay and silt	For most crops, a soil texture of 7–27% clay, 28–50% silt, and 23–52% sand is thought to be optimal. (Bouyoucos hydrometer method and International pipette method)
Bulky density	Mg m ⁻³ or Gram cm ⁻³	1.33–1.35 g cm ⁻³ (Direct and indirect methods)
Penetration resistance	MegaPascal (MPa); N m ⁻² (cone index N cm ⁻²)	N/A Cone penetrometer
Aggregate stability	Mean weight diameter (mm); Geometric mean diameter (mm)	N/A (Wet sieving and dry sieving method)
Water holding capacity	mm m ⁻¹ depth of soil	Crops specific (Pressure plate and membrane apparatus)
Infiltration rate	mm hour ⁻¹	N/A (Ring infiltrometer)
Depth of hardpan	Indicated as depth from the surface at which hardpan observe	Based on the effective root zone depth and characteristics of plant (Determined by compaction of soil at different layers)

Depth of water table	Depth from the surface in meters	N/A (Paizometer and open well)
Porosity	Percentage%	50% of the total soil volume (Mercury intrusion porosimetry; Image analysis and soil micromorphology)
Erosive potential	Mg ha ⁻¹ soil loss year ⁻¹	≤ 11 Mg ha ⁻¹ soil loss/year (permissible limit) (Universal soil loss equation)
Soil Structure	Expressed as types (Platy, prismatic, blocky and spheroidal), class (Very fine, fine/thin, medium, coarse/thick and very coarse) and grade (structureless, weak, moderate and strong)	N/A
Soil crust	Qualitative property indicated by either types of crust or by surface hardness measured by cone penitrometer	Soil should be crust free as all crust has adverse from cultivation point of view except soil biological crust in some cases (Optical and scanning electron microscopy)
PH	In scale of 1–14	Neutral (6.7–7.3) pH for most of the crops and soil functioning is considered as ideal
Electrical conductivity	dS m ⁻¹	N/A (Saturation soil extract or soil- water suspension)
Minor nutrients	2.0 mg kg ⁻¹ soil Zinc (Zn)	N/A
Manganese Copper zinc Boron	0.6 mg kg ⁻¹ soil Copper (Cu) 0.2 mg kg ⁻¹ soil Boron (B) mg kg ⁻¹ 0.5 mg kg ⁻¹ soil	
Urease Enzyme	N/A	Soil incubation in tri(hydroxymethyl) aminomethane buffer
Microbial biomass carbon	(µg microbial biomass carbon g ⁻¹ soil)	N/A (Fumigation method)

Results

Important conclusions on conservation tillage's effects on soil preservation and carbon sequestration can be gathered from a survey of the research.

Conservation tillage techniques improve carbon sequestration in agricultural soils

It is well known that conservation tillage, a commonly used agricultural technique, protects soil resources by increasing soil organic carbon (SOC) (Zhu *et al.*, 2022).

(Lal *et al.*, 1997) one of the information collected the two main ways that conservation tillage sequesters carbon are by deep storing SOC in the subsoil horizons and promoting micro-aggregation. Increased biomass output through conservation tillage and related agricultural methods (e.g., soil fertility enhancement, improved crops and species, cover crops and fallowing, improved pastures and deep-rooted crops) are also beneficial.

(Haddaway *et al.*, 2017) reported conservation tillage, such as decreased fuel usage and erosion, there are also negative effects, such as N₂O etc.

(Hussain *et al.*, 2021) It has been suggested that the main goal of conservation agriculture is to enhance soil health and plant development without causing adverse effects on the environment.

(Francaviglia *et al.*, 2023) studied By improving soil organic carbon (SOC) sequestration in soils and its associated co-benefits, sustainable agricultural practices—such as reducing tillage, cultivating cover crops, and in place crop residue retention measures—have been suggested as low-cost solutions that can address land degradation, food security, and climate change mitigation and adaptation. Accordingly, a great deal of research has shown that conservation agriculture (CA) enhances the biological, chemical, and physical features of soil, all of which are essential for preserving soil health and improving the adaptability of agroecosystems to climate change.

as numerous studies have repeatedly shown. The fundamental reason for this is the decrease in soil disturbance, which promotes a favorable environment for carbon storage and aids in maintaining organic matter levels.

Conservation tillage can be a practical way to store carbon in the soil and minimize the effects of climate change

(Deng *et al.*, 2022) one of the information collected thus, conservation tillage improves climatic resilience and minimizes the effects of climate change on agriculture.

(Rahman *et al.*, 2021) suggested Intensive soil tillage and crop residue removal in conventional agricultural systems may have a greater severe impact on the environment.

(Alhassan *et al.*, 2021) reported shown that NTS, in particular, enhanced soil water content and decreased soil temperature through conservation tillage techniques.

(Yao *et al.*, 2023) investigated if conservation tillage techniques may decrease the impact of climate change on soil CO₂ emissions from arid farms.

Based on our research, zero tillage may be a major factor in reducing greenhouse gas emissions from soils and aiding in the fight against climate change.

Moreover, conservation tillage is essential to preserving soil health, according to the examination of soil preservation indicators.

Under conservation tillage systems, studies regularly show benefits in soil structure, moisture retention, and nutrient levels. Improved water infiltration and decreased erosion are noted, which over time will lead to better soil maintenance. The findings support the sustainability of agricultural ecosystems by confirming that conservation tillage practices have a positive impact on a number of soil quality factors.

Conservation tillage reduce greenhouse gas intensity in organic farming

(D.E *et al.*, 2001) showed that tillage has an organic soil-saving effect that can lower greenhouse gas emissions and future farm fuel usage while also preserving energy for increased profit.

(Rahman *et al.*, 2021) suggests MT and ZT practices to reduce adverse environmental impacts in Bangladeshi wheat agriculture, as the results support CTS. When comparing the techniques, the MT method—which keeps the crop residue (20 cm) and applies CA principles—is more suited for Bangladesh's wheat agriculture, both for CSA and SI. This is because it can enhance SOC formation while preventing water loss and greenhouse gas emissions without compromising output.

(Valujeva et al., 2022) There have been suggestions for reduced tillage and alternative crops to lower greenhouse gas emissions from agricultural soils.

(Gryze et al., 2010) reported organic farming, winter cover crops, and conservation tillage have all been suggested as strategies to lower soil greenhouse gas emissions from agriculture.

(Khresat et al., 2016) showed that conservation agriculture practices lower the greenhouse gas emissions of farming systems. (Khan *et al.*, 2023) studied Carbon sequestration can reduced the green house gas emission.

Discussion

The findings are consistent with the idea that conservation tillage is a viable strategy for reducing the effects of climate change and maintaining soil health. Increased soil organic carbon content is a result of both crop residues remaining on the field surface as well as decreased soil disturbance. This thus helps with carbon sequestration, resolving the issue of greenhouse gas emissions. The benefits for preserving soil are examined in relation to better water management and erosion prevention. Conservation tillage techniques reduce soil erosion by keeping surface leftovers in place, which serve as a protective layer. Sustainable soil management necessitates improved water infiltration and moisture retention, both of which increase resistance to drought.

It is important to recognize potential challenges and limitations linked to conservation tillage. In some situations, localized variables like crop rotation techniques, soil composition, and climate might affect how effective conservation tillage is. Farmers may face initial difficulties if these practices are adopted since they may call for changes to machinery and management.

The debate and overall findings highlight the significance of conservation tillage as a sustainable farming method for soil protection and carbon sequestration. The results provide insightful information to guide future investigations as well as promote the adoption of strategies that improve agricultural systems' long-term sustainability.

Conclusions and future perspectives

The best way to combat the negative consequences of climate change on agriculture, a sector that is extremely sensitive to changing weather patterns, is to manage natural resources carefully. Transferring atmospheric CO₂ into the soil through a process known as "soil C sequestration" is a mutually beneficial strategy that addresses both climate adaptation and mitigation. Plant photosynthesis is the main process that converts atmospheric CO₂ into soil, and it entails defending the soil's carbon-based pools from soil microbial populations that would otherwise release the carbon back into the atmosphere. The no-till farming method is regarded as an efficient way to restore soil and absorb atmospheric carbon since it maintains ecosystems and soil health. In addition to improving the efficiency of water and fertilizer use, zero- or no-tillage when combined with keeping crop residue in the field or using it as mulch helps sequester a sizable amount of atmospheric CO₂. Crop rotation has the potential to improve soil health and sequester carbon under a conservation agriculture system by accelerating SOC accumulation rates at different soil levels. The majority of agricultural management methods that support carbon sequestration also enhance soil fertility, increase soil aggregate stability, retain water better, and guarantee food security. However, taking action shouldn't be contingent on having a thorough understanding of soil C and the sequestration capacity. Numerous techniques to improve the sequestration of atmospheric C have recently been presented by diverse research projects on various agricultural management methods. The adaptation of conservation tillage practices is comparatively more effective than several other options for atmospheric drawdown, and it may be adapted soon. Risks involved in this system are low, and there are several established advantages to enhancing soil quality and sequestering C.

Acknowledgment: I am very thankful to China Scholarship Council. This research supported by Chinese government scholarship.

Conflict of interest: The author declare there is no conflict of interest.

Funding: No financial support was given to the author(s) for the research, writing, or publication of this paper.

Authors contribution: Conceived and designed the research project: Waqar Ahmad Khan & Gang Wang , Performed the review paper: Gang Wang , Analyzed, Waqar Ahmad Khan Wrote this paper.

Data availability: N/A

References

- Abiven, S.; Menasseri, S.; Chenu, C. The effects of organic inputs over time on soil aggregate stability—A literature analysis. *Soil Biol. Biochem.* 2009, 41, 1–12.
- Allmaras, R.R.; Linden, D.R.; Clapp, C. Corn-residue transformations into root and soil carbon as related to nitrogen, tillage, and stover management. *Soil Sci. Soc. Am. J.* 2004, 68, 1366–1375.
- Alhassan, A. R. M., Yang, C., Ma, W., & Li, G. (2021). Influence of conservation tillage on Greenhouse gas fluxes and crop productivity in spring-wheat agroecosystems on the Loess Plateau of China. *PeerJ*, 9, e11064.
- Assessment, F. G. F. R. (2015). How are the world's forests changing. Food and the Agricultural Organization: Rome, Italy, 47.
- Anikwe, M. A. N., & Ubochi, J. N. (2007). Short-term changes in soil properties under tillage systems and their effect on sweet potato (*Ipomea batatas* L.) growth and yield in an Ultisol in south-eastern Nigeria. *Soil Research*, 45(5), 351-358.
- Allmaras, R. R., Rickman, R. W., Ekin, L. G., & Kimball, B. A. (1977). Chiseling influences on soil hydraulic properties. *Soil Science Society of America Journal*, 41(4), 796-803.
- Busari, M.A.; Kukal, S.S.; Kaur, A.; Bhatt, R.; Dulazi, A.A. Conservation tillage impacts on soil, crop and the environment. *Int. Soil Water Conserv. Res.* 2015, 3, 119–129.
- Beare, M.; Hendrix, P.; Coleman, D. Water-stable aggregates and organic matter fractions in conventional-and no-tillage soils. *Soil Sci. Soc. Am. J.* 1994, 58, 777–786.
- Bhattacharyya, R.; Das, T.; Sudhishri, S.; Dudwal, B.; Sharma, A.; Bhatia, A.; Singh, G. Conservation agriculture effects on soil organic carbon accumulation and crop productivity under a rice–wheat cropping system in the western Indo-Gangetic Plains. *Eur. J. Agron.* 2015, 70, 11–21.
- Bossuyt, H.; Six, J.; Hendrix, P.F. Aggregate-protected carbon in no-tillage and conventional tillage agroecosystems using carbon-14 labeled plant residue. *Soil Sci. Soc. Am. J.* 2002, 66, 1965–1973.
- Beare, M.H.; Hendrix, P.; Cabrera, M.; Coleman, D. Aggregate-protected and unprotected organic matter pools in conventional and no-tillage soils. *Soil Sci. Soc. Am. J.* 1994, 58, 787–795.
- Blaikie, P., & Brookfield, H. (Eds.). (2015). *Land degradation and society*. Routledge.
- Blum, W. E., & Swaran, H. (2004). Soils for sustaining global food production. *Journal of food science*, 69(2), crh37-cr42.
- Beinroth, F. H., Eswaran, H., Reich, P. F., & Van Den Berg, E. (1994). Land related stresses in agroecosystems. *Stressed ecosystems and sustainable agriculture*.
- Bhatt, R., & Khera, K. L. (2006). Effect of tillage and mode of straw mulch application on soil erosion in the submontaneous tract of Punjab, India. *Soil and Tillage Research*, 88(1-2), 107-115.
- Benjamin, J. G. (1993). Tillage effects on near-surface soil hydraulic properties. *Soil and Tillage Research*, 26(4), 277-288.

- Busari, M. A., & Salako, F. K. (2012). Effect of tillage and poultry manure application on soil infiltration rate and maize root growth in a sandy Alfisol. *Agro-Science*, 11(2), 24-31.
- Babu, S., Singh, R., Avasthe, R., Rathore, S. S., Kumar, S., Das, A., ... & Singh, V. K. (2023). Conservation tillage and diversified cropping enhance system productivity and eco-efficiency and reduce greenhouse gas intensity in organic farming. *Frontiers in Sustainable Food Systems*, 7, 1114617.
- Butorac, A. (2017). Conservation tillage in eastern Europe. In *Conservation tillage in temperate agroecosystems* (pp. 357-374). CRC Press.
- Corsi, S., Friedrich, T., Kassam, A., Pisante, M., & de Moraes Sà, J. C. (2012). Soil organic carbon accumulation and greenhouse gas emission reductions from conservation agriculture: A literature review, integrated crop management (101 pp.). Vol. 16. Rome: AGP/FAO.
- Conservation Technology Information Center. (1992). National crop residue management survey. NACD's Conservation Technology Information Center.
- Campbell, C.; Selles, F.; Lafond, G.; Zentner, R. Adopting zero tillage management: Impact on soil C and N under long-term crop rotations in a thin Black Chernozem. *Can. J. Soil Sci.* 2001, 81, 139–148.
- Carter, M.R. Soil quality for sustainable land management: Organic matter and aggregation interactions that maintain soil functions. *Agron. J.* 2002, 94, 38–47.
- Charreau, C. (1972). Problèmes posés par l'utilisation agricole des sols tropicaux par des cultures annuelles.
- Chisholm, A., & Dumsday, R. (Eds.). (1987). *Land degradation: problems and policies* (No. 18). Cambridge University Press.
- Carter, M. R. (2005). Conservation tillage. *Encyclopedia of Soils in the Environment*, 4, 306-311.
- Dumanski, J.; Peiretti, R.; Benites, J.; McGarry, D.; Pieri, C. The paradigm of conservation agriculture. *Proc. World Assoc. Soil Water Conserv.* 2006, 1, 58–64.
- De Gouvello, C. Brazil Low-carbon Country Case Study; The World Bank Group; The International Bank for Reconstruction and Development; The World Bank: Washington, DC, USA, 2010.
- Deng, X.; Yang, Q.; Zhang, D.; Dong, S. Application of Conservation Tillage in China: A Method to Improve Climate Resilience. *Agronomy* **2022**, 12, 1575. <https://doi.org/10.3390/agronomy12071575>
- de Magalhães, M.M.; Lima, D.L. Low-Carbon Agriculture in Brazil: The Environmental and Trade Impact of Current Farm Policies, Issue Paper No. 54; International Centre for Trade and Sustainable Development: Geneva, Switzerland, 2014.
- Das, A., Basavaraj, S., Layek, J., Gandhiji Idapuganti, R., Lal, R., Rangappa, K., et al. (2019). Can conservation tillage and residue management enhance energy use efficiency and sustainability of rice-vegetable pea systems in the Eastern Himalayas? *Arch. Agron. Soil Sci.* 66, 830–846. doi: 10.1080/03650340.2019.1639157
- Ella, V. B., Reyes, M. R., Mercado Jr, A., Adrian, A., & Padre, R. (2016). Conservation agriculture increases soil organic carbon and residual water content in upland crop production systems. *Eurasian Journal of Soil Science*, 5(1), 24-29.
- Eswaran, H., Reich, P., & Beinroth, F. (1997). Global distribution of soils with acidity. P 159-164. *Plantsoil interactions at low pH*. Brazilian soil sci. Society. Campinas/vicosa, Brazil.
- Eswaran, H., Lal, R., & Reich, P. F. (2001). Land degradation: an overview. Responses to Land Degradation. In *Proc. 2nd International Conference on Land Degradation and Desertification*. Khon Kaen, Thailand, edited by E. Bridges, I. Hannam, L. Oldeman, F. Penning de Vries, S. Scherr, and S. Sompatpanit. New Delhi: Oxford Press.
- Eriksson, J., Håkansson, I., & Danfors, B. (1974). Jordpackning-markstruktur-gröda [The effect of soil compaction on soil structure and crop yields]. In Rep. 354 (pp. 1-82). Swedish Institute of Agricultural Engineering.

- Francaviglia, R.; Almagro, M.; Vicente-Vicente, J.L. Conservation Agriculture and Soil Organic Carbon: Principles, Processes, Practices and Policy Options. *Soil Syst.* **2023**, *7*, 17. <https://doi.org/10.3390/soilsystems7010017>
- Giller, K.E.; Witter, E.; Corbeels, M.; Titttonell, P. Conservation agriculture and smallholder farming in Africa: The heretics' view. *Field Crop. Res.* **2009**, *114*, 23–34.
- Gregorich, E.; Monreal, C.; Ellert, B. Turnover of soil organic matter and storage of corn residue carbon estimated from natural ¹³C abundance. *Can. J. Soil Sci.* **1995**, *75*, 161–167.
- Gale, W.; Cambardella, C.; Bailey, T. Root-derived carbon and the formation and stabilization of aggregates. *Soil Sci. Soc. Am. J.* **2000**, *64*, 201–207.
- Gryze, S. D., Wolf, A., Kaffka, S. R., Mitchell, J., Rolston, D. E., Temple, S. R., ... & Six, J. (2010). Simulating greenhouse gas budgets of four California cropping systems under conventional and alternative management. *Ecological applications*, *20*(7), 1805-1819.
- Gebara, M.F.; Thuault, A. GHG Mitigation in Brazil's Land Use Sector: An Introduction to the Current National Policy Landscape; WRI:Washington, DC, USA, 2013.
- Gupta, D.; Bhatia, A.; Kumar, A.; Chakrabarti, B.; Jain, N.; Pathak, H. Global warming potential of rice (*Oryza sativa*)-wheat (*Triticum aestivum*) cropping system of the Indo-Gangetic Plains. *Indian J. Agric. Sci* **2015**, *85*, 807–816.
- González-Sánchez, E.; Ordóñez-Fernández, R.; Carbonell-Bojollo, R.; Veroz-González, O.; Gil-Ribes, J. Meta-analysis on atmospheric carbon capture in Spain through the use of conservation agriculture. *Soil Tillage Res.* **2012**, *122*, 52–60.
- Garibaldi, L. (2012). The FAO global capture production database: a six-decade effort to catch the trend. *Marine Policy*, *36*(3), 760-768.
- Gregorich, E.; Drury, C.; Baldock, J. Changes in soil carbon under long-term maize in monoculture and legume-based rotation. *Can. J. Soil Sci.* **2001**, *81*, 21–31.
- Himes, F. Nitrogen, sulfur, and phosphorus and the sequestering of carbon. In *Soil Processes and the Carbon Cycle*; CRC Press: Boca Raton, FL, USA, 2018; pp. 315–319.
- Hille, D.; Rosenzweig, C. The role of soils in climate change. In *Handbook of Climate Change and Agroecosystems: Impacts, Adaptation, and Mitigation*; Hille, D., Rosenzweig, C., Eds.; ICP Series on Climate Change Impacts, Adaptation, and Mitigation; Imperial College Press: Singapore, 2011; Volume 1, pp. 9–20.
- Hedlund, A., Witter, E., & An, B. X. (2003). Assessment of N, P and K management by nutrient balances and flows on peri-urban smallholder farms in southern Vietnam. *European Journal of Agronomy*, *20*(1-2), 71-87.
- Hussain, S., Hussain, S., Guo, R., Sarwar, M., Ren, X., Krstic, D., ... & El-Esawi, M. A. (2021). Carbon sequestration to avoid soil degradation: A review on the role of conservation tillage. *Plants*, *10*(10), 2001.
- Haddaway, N. R., Hedlund, K., Jackson, L. E., Kätterer, T., Lugato, E., Thomsen, I. K., ... & Isberg, P. E. (2017). How does tillage intensity affect soil organic carbon? A systematic review. *Environmental Evidence*, *6*(1), 1-48.
- Haile, S.G.; Nair, P.R.; Nair, V.D. Carbon storage of different soil-size fractions in Florida silvopastoral systems. *J. Environ. Qual.* **2008**, *37*, 1789–1797.
- Ivanova, A. (2011). Carbon Sequestration.
- Jain, N.; Dubey, R.; Dubey, D.; Singh, J.; Khanna, M.; Pathak, H.; Bhatia, A. Mitigation of greenhouse gas emission with system of rice intensification in the Indo-Gangetic Plains. *Paddy Water Environ.* **2014**, *12*, 355–363.
- Jain, N.; Dubey, R.; Dubey, D.; Singh, J.; Khanna, M.; Pathak, H.; Bhatia, A. Mitigation of greenhouse gas emission with system of rice intensification in the Indo-Gangetic Plains. *Paddy Water Environ.* **2014**, *12*, 355–363.

- Jacinthe, P.; Lal, R.; Kimble, J. Effects of wheat residue fertilization on accumulation and biochemical attributes of organic carbon in a central Ohio Luvisol. *Soil Sci.* 2002, 167, 750–758.
- Jacobs, A., Rauber, R., & Ludwig, B. (2009). Impact of reduced tillage on carbon and nitrogen storage of two Haplic Luvisols after 40 years. *Soil and Tillage Research*, 102(1), 158-164.
- Jastrow, J.; Miller, R. Methods for assessing the effects of biota on soil structure. *Agric. Ecosyst. Environ.* 1991, 34, 279–303.
- Khan Waqar Ahmad, & Gang Wang. (2023). Evaluating the Crucial Relationships between Soil Health and Climate Change. *Journal of Environmental Impact and Management Policy (JEIMP)* ISSN:2799-113X, 4(01), 8–21. <https://doi.org/10.55529/jeimp.41.8.21>
- Kayombo, B., & Lal, R. (1994). Responses of tropical crops to soil compaction. In *Developments in agricultural engineering* (Vol. 11, pp. 287-316). Elsevier.
- Khresat, S. (2016, April). Practicing Conservation Agriculture to mitigate and adapt to Climate Change in Jordan. In *EGU General Assembly Conference Abstracts* (pp. EPSC2016-685).
- Karami, A.; Homaei, M.; Afzalnia, S.; Ruhipour, H.; Basirat, S. Organic resource management: Impacts on soil aggregate stability and other soil physico-chemical properties. *Agric. Ecosyst. Environ.* 2012, 148, 22–28.
- Kahlon, M.S.; Lal, R.; Ann-Varughese, M. Twenty two years of tillage and mulching impacts on soil physical characteristics and carbon sequestration in Central Ohio. *Soil Tillage Res.* 2013, 126, 151–158.
- Kou, T.; Zhu, P.; Huang, S.; Peng, X.; Song, Z.; Deng, A.; Gao, H.; Peng, C.; Zhang, W. Effects of long-term cropping regimes on soil carbon sequestration and aggregate composition in rainfed farmland of Northeast China. *Soil Tillage Res.* 2012, 118, 132–138.
- Kertész, Á. (2009). The global problem of land degradation and desertification. *Hungarian Geographical Bulletin*, 58(1), 19-31.
- Kargas, G., Kerkides, P., & Poulouvassilis, A. (2012). Infiltration of rain water in semi-arid areas under three land surface treatments. *Soil and Tillage Research*, 120, 15-24.
- Kemper, W. D., Trout, T. J., Segeren, A., & Bullock, M. (1987). Worms and water. *Journal of Soil and Water Conservation*, 42(6), 401-404.
- Lal, R. (1990). *Soil erosion in the tropics: principles and management*. McGraw-Hill Inc..
- Li, L. L., Huang, G. B., Zhang, R. Z., Jin, X. J., Li, G. D., & Chan, K. Y. (2005). Effects of conservation tillage on soil water regimes in rainfed areas. *Acta Ecologica Sinica*, 25(9), 2326-2332.
- Lal, R. Soil carbon sequestration impacts on global climate change and food security. *Science* 2004, 304, 1623–1627.
- Lal, R. (1996). Deforestation and land-use effects on soil degradation and rehabilitation in western Nigeria. I. Soil physical and hydrological properties. *Land degradation & development*, 7(1), 19-45.
- Lal, R. Soil management and restoration for C sequestration to mitigate the accelerated greenhouse effect. *Prog. Environ. Sci.* 1999,1, 307–326.
- Lal, R.; Kimble, J. Conservation tillage for carbon sequestration. *Nutr. Cycl. Agroecosyst.* 1997, 49, 243–253.
- Lal, R. Challenges and opportunities in soil organic matter research. *Eur. J. Soil Sci.* 2009, 60, 158–169.
- Lal, R. Carbon sequestration in dryland ecosystems. *Environ. Manag.* 2004, 33, 528–544.
- Lal, R. Residue management, conservation tillage and soil restoration for mitigating greenhouse effect by CO₂-enrichment. *Soil Tillage Res.* 1997, 43, 81 107.
- Lal, R. Digging deeper: A holistic perspective of factors affecting soil organic carbon sequestration in agroecosystems. *Glob.Chang. Biol.* 2018, 24, 3285–3301.
- Lal, R. (2020). *Soil Quality and Sustainability. Methods for Assessment of Soil Degradation*, 17.

- Lal, R. Soil carbon sequestration to mitigate climate change. *Geoderma* 2004, 123, 1–22.
- Lavelle, P.; Pashanasi, B. Soil macrofauna and land management in Peruvian Amazonia (Yurimaguas, Loreto). *Pedobiologia* 1989, 33, 283–291.
- Lorenz, K.; Lal, R. The depth distribution of soil organic carbon in relation to land use and management and the potential of carbon sequestration in subsoil horizons. *Adv. Agron.* 2005, 88, 35–66.
- Lal, R. (1987). Response of maize (*Zea mays*) and cassava (*Manihot esculenta*) to removal of surface soil from an Alfisol in Nigeria. *International Journal of Tropical Agriculture*, 5(2), 77–92.
- Lal, R. (1995). Erosion-crop productivity relationships for soils of Africa. *Soil Science Society of America Journal*, 59(3), 661–667.
- Murungu, F.; Chidzuza, C.; Muchaonyerwa, P.; Mkeni, P. Mulch effects on soil moisture and nitrogen, weed growth and irrigated maize productivity in a warm-Temp. climate of South Africa. *Soil Tillage Res.* 2011, 112, 58–65.
- Mbagwu, J. S. C., Lal, R., & Scott, T. W. (1984). Effects of desurfacing of Alfisols and Ultisols in southern Nigeria: I. Crop performance. *Soil Science Society of America Journal*, 48(4), 828–833.
- McVay, K. A., Budde, J. A., Fabrizzi, K., Mikha, M. M., Rice, C. W., Schlegel, A. J., ... & Thompson, C. (2006). Management effects on soil physical properties in long-term tillage studies in Kansas. *Soil Science Society of America Journal*, 70(2), 434–438.
- Martínez, E., Fuentes, J. P., Silva, P., Valle, S., & Acevedo, E. (2008). Soil physical properties and wheat root growth as affected by no-tillage and conventional tillage systems in a Mediterranean environment of Chile. *Soil and Tillage Research*, 99(2), 232–244.
- Navarro-Pedreño, J.; Almendro-Candel, M.B.; Zorpas, A.A. The increase of soil organic matter reduces global warming, myth or reality? *Science* 2021, 3, 18.
- Mangalassery S, Sjögersten S, Sparkes DL, Sturrock CJ, Craigon J, Mooney SJ. To what extent can zero tillage lead to a reduction in greenhouse gas emissions from temperate soils? *Sci Rep.* 2014 Apr 4;4:4586. doi: 10.1038/srep04586. PMID: 24699273; PMCID: PMC3975454.
- Obalum, S.; Obi, M. Physical properties of a sandy loam Ultisol as affected by tillage-mulch management practices and cropping systems. *Soil Tillage Res.* 2010, 108, 30–36.
- Oldeman, L. R. (1994). An international methodology for an assessment of soil degradation, land georeferenced soils and terrain database. RAPA Publication (FAO).
- Oldeman, L. R. (1992). Global extent of soil degradation. In Bi-annual report 1991–1992/ISRIC (pp. 19–36). ISRIC.
- Olson, K.R.; Al-Kaisi, M.; Lal, R.; Cihacek, L. Impact of soil erosion on soil organic carbon stocks. *J. Soil Water Conserv.* 2016, 71, 61A–67A.
- Puget, P.; Lal, R.; Izaurrealde, C.; Post, M.; Owens, L. Stock and distribution of total and corn-derived soil organic carbon in aggregate and primary particle fractions for different land use and soil management practices. *Soil Sci.* 2005, 170, 256–279.
- Palm, C.; Blanco-Canqui, H.; DeClerck, F.; Gatere, L.; Grace, P. Conservation agriculture and ecosystem services: An overview. *Agric. Ecosyst. Environ.* 2014, 187, 87–105.
- Pisante, M.; Stagnari, F.; Acutis, M.; Bindi, M.; Brilli, L.; Di Stefano, V.; Carozzi, M. Conservation agriculture and climate change. In *Conservation Agriculture*; Springer: Berlin/Heidelberg, Germany, 2015; pp. 579–620.
- Pisante, M.; Stagnari, F.; Grant, C.A. Agricultural innovations for sustainable crop production intensification. *Ital. J. Agron.* 2012, 7, e40.
- Prasad, J.; Rao, C.S.; Srinivas, K.; Jyothi, C.N.; Venkateswarlu, B.; Ramachandrapa, B.; Dhanapal, G.; Ravichandra, K.; Mishra, P. Effect of ten years of reduced tillage and recycling of organic matter on crop

- yields, soil organic carbon and its fractions in Alfisols of semi arid tropics of southern India. *Soil Tillage Res.* 2016, 156, 131–139.
- Pagliai, M., Vignozzi, N., & Pellegrini, S. (2004). Soil structure and the effect of management practices. *Soil and tillage research*, 79(2), 131-143.
- Pikul Jr, J. L., & Aase, J. K. (1995). Infiltration and soil properties as affected by annual cropping in the northern Great Plains. *Agronomy Journal*, 87(4), 656-662.
- Rashidi, M.; Keshavarzpour, F. Effect of different tillage methods on grain yield and yield components of maize (*Zea mays* L.). *Int.J. Agric. Biol* 2007, 9, 274–277.
- Raj, R., Das, T. K., Pankaj, S., Banerjee, T., Ghosh, A., Bhattacharyya, R., et al. (2022). Co-implementation of conservation tillage and herbicides reduces weed and nematode infestation and enhances the productivity of direct-seeded rice in North-western Indo-Gangetic Plains. *Front. Sustain. Food Syst.* 6, 1017013. doi: 10.3389/fsufs.2022.1017013
- Rashidi, M.; Keshavarzpour, F. Effect of different tillage methods on soil physical properties and crop yield of melon (*Cucumis melo*). *ARPN J. Agric. Biol. Sci.* 2008, 3, 41–46.
- Rashidi, M.; Gholami, M.; Abbassi, S. Effect of different tillage methods on yield and yield components of tomato (*Lycopersicon esculentum*). *ARPN J. Agric. Biol. Sci.* 2006, 5, 26–30.
- Rahman, M. M., Aravindakshan, S., Hoque, M. A., Rahman, M. A., Gulandaz, M. A., Rahman, J., & Islam, M. T. (2021). Conservation tillage (CT) for climate-smart sustainable intensification: Assessing the impact of CT on soil organic carbon accumulation, greenhouse gas emission and water footprint of wheat cultivation in Bangladesh. *Environmental and Sustainability Indicators*, 10, 100106.
- Ruppenthal, M. (1995). Soil conservation in Andean crop** systems: soil erosion and crop productivity in traditional and forage-legume based cassava crop** systems in the south Colombian Andes.
- Srinivasarao, C.; Vittal, K.; Venkateswarlu, B.; Wani, S.; Sahrawat, K.; Marimuthu, S.; Kundu, S. Carbon stocks in different soil types under diverse rainfed production systems in tropical India. *Commun. Soil Sci. Plant Anal.* 2009, 40, 2338–2356.
- Shrestha, B.; McConkey, B.; Smith, W.; Desjardins, R.; Campbell, C.; Grant, B.; Miller, P. Effects of crop rotation, crop type and tillage on soil organic carbon in a semiarid climate. *Can. J. Soil Sci.* 2013, 93, 137–146.
- Scherr, S. J. (1999). Soil degradation: a threat to develo**-country food security by 2020? (Vol. 27). *Intl Food Policy Res Inst.*
- Somasundaram, J.; Sinha, N.; Dalal, R.C.; Lal, R.; Mohanty, M.; Naorem, A.; Hati, K.; Chaudhary, R.; Biswas, A.; Patra, A. No-till farming and conservation agriculture in South Asia—issues, challenges, prospects and benefits. *Crit. Rev. Plant Sci.* 2020, 39,236–279.
- Shahane, A. A., & Shivay, Y. S. (2021). Soil health and its improvement through novel agronomic and innovative approaches. *Frontiers in Agronomy*, 3, 680456.
- Six, J.; Elliott, E.; Paustian, K. Aggregate and soil organic matter dynamics under conventional and no-tillage systems. *Soil Sci. Soc. Am. J.* 1999, 63, 1350–1358.
- Srinivasarao, C.; Venkateswarlu, B.; Lal, R.; Singh, A.K.; Kundu, S.; Vittal, K.P.R.; Ramachandrappa, B.K.; Gajanan, G.N. Longterm effects of crop residues and fertility management on carbon sequestration and agronomic productivity of groundnut–finger millet rotation on an Alfisol in southern India. *Int. J. Agric. Sustain.* 2012, 10, 230–244.
- Six, J.; Paustian, K.; Elliott, E.T.; Combrink, C. Soil structure and organic matter I. Distribution of aggregate-size classes and aggregate-associated carbon. *Soil Sci. Soc. Am. J.* 2000, 64, 681–689.
- Stocking, M. A. (2003). Tropical soils and food security: the next 50 years. *Science*, 302(5649), 1356-1359.

- Shrestha, B.; McConkey, B.; Smith, W.; Desjardins, R.; Campbell, C.; Grant, B.; Miller, P. Effects of crop rotation, crop type and tillage on soil organic carbon in a semiarid climate. *Can. J. Soil Sci.* 2013, 93, 137–146.
- Schumacher, T. E., Lindstrom, M. J., Mokma, D. L., & Nelson, W. W. (1994). Corn yield: Erosion relationships of representative loess and till soils in the North Central United States. *Journal of Soil and Water Conservation*, 49(1), 77-81.
- Six, J.; Conant, R.T.; Paul, E.A.; Paustian, K. Stabilization mechanisms of soil organic matter: Implications for C-saturation of soils. *Plant Soil* 2002, 241, 155–176.
- Silburn, D. M., Freebairn, D. M., & Rattray, D. J. (2007). Tillage and the environment in sub-tropical Australia—Tradeoffs and challenges. *Soil and Tillage Research*, 97(2), 306-317.
- Su, Z., Zhang, J., Wu, W., Cai, D., Lv, J., Jiang, G., ... & Gabriels, D. (2007). Effects of conservation tillage practices on winter wheat water-use efficiency and crop yield on the Loess Plateau, China. *Agricultural Water Management*, 87(3), 307-314.
- Shukla, M. K., Lal, R., Owens, L. B., & Unkefer, P. (2003). Land use and management impacts on structure and infiltration characteristics of soils in the North Appalachian region of Ohio. *Soil Science*, 168(3), 167-177.
- Tisdall, J.M.; Oades, J.M. Organic matter and water-stable aggregates in soils. *J. Soil Sci.* 1982, 33, 141–163.
- Thierfelder, C.; Wall, P.C. Effects of conservation agriculture techniques on infiltration and soil water content in Zambia and Zimbabwe. *Soil Tillage Res.* 2009, 105, 217–227.
- Tejada, M.; Garcia, C.; Gonzalez, J.; Hernandez, M. Organic amendment based on fresh and composted beet vinasse: Influence on soil properties and wheat yield. *Soil Sci. Soc. Am. J.* 2006, 70, 900–908.
- Toor, G. S., Yang, Y. Y., Das, S., Dorsey, S., & Felton, G. (2021). Soil health in agricultural ecosystems: current status and future perspectives. *Advances in Agronomy*, 168, 157-201.
- Verhulst, N.; Govaerts, B.; Verachtert, E.; Castellanos-Navarrete, A.; Mezzalama, M.; Wall, P.; Deckers, J.; Sayre, K.D. Conservation agriculture, improving soil quality for sustainable production systems. In *Advances in Soil Science: Food Security and Soil Quality*, CRC Press: Boca Raton, FL, USA, 2010; pp. 137–208.
- Vicente-Vicente, J. L., García-Ruiz, R., Francaviglia, R., Aguilera, E., & Smith, P. (2016). Soil carbon sequestration rates under Mediterranean woody crops using recommended management practices: A meta-analysis. *Agriculture, Ecosystems & Environment*, 235, 204-214.
- Valujeva, K., Pilecka-Ulcugaceva, J., Skiste, O., Liepa, S., Lagzdins, A., & Grinfelde, I. (2022). Soil tillage and agricultural crops affect greenhouse gas emissions from Cambic Calcisol in a temperate climate. *Acta Agriculturae Scandinavica, Section B—Soil & Plant Science*, 72(1), 835-846.
- Wanniarachchi, S.; Voroney, R.; Vyn, T.; Beyaert, R.; MacKenzie, A. Tillage effects on the dynamics of total and corn-residue-derived soil organic matter in two southern Ontario soils. *Can. J. Soil Sci.* 1999, 79, 473–480.
- Wani, S.A.; Mehraj-Ud-din, K.; Bashir, Z.; Kousar, S.; Rasool, F.; Zargar, M. Carbon Sequestration Potential of Soils—A Review. *Carbon Sequestration Potential Soils* 2016, 5, 9500–9600.
- Wanniarachchi, S.; Voroney, R.; Vyn, T.; Beyaert, R.; MacKenzie, A. Tillage effects on the dynamics of total and corn-residue-derived soil organic matter in two southern Ontario soils. *Can. J. Soil Sci.* 1999, 79, 473–480.
- Wani, S.A.; Mehraj-Ud-din, K.; Bashir, Z.; Kousar, S.; Rasool, F.; Zargar, M. Carbon Sequestration Potential of Soils—A Review. *Carbon Sequestration Potential Soils* 2016, 5, 9500–9600.
- Yu, H.; Ding, W.; Chen, Z.; Zhang, H.; Luo, J.; Bolan, N. Accumulation of organic C components in soil and aggregates. *Sci. Rep.* 2015, 5, 1–12.
- Yao, Y., Li, G., Lu, Y., & Liu, S. (2023). Modelling the impact of climate change and tillage practices on soil CO₂ emissions from dry farmland in the Loess Plateau of China. *Ecological Modelling*, 478, 110276.
- Yang, X.; Drury, C.; Reynolds, W.; Tan, C. Impacts of long-term and recently imposed tillage practices on the vertical distribution of soil organic carbon. *Soil Tillage Res.* 2008, 100, 120–124.

- Zhang, M.; Cheng, G.; Feng, H.; Sun, B.; Zhao, Y.; Chen, H.; Chen, J.; Dyck, M.; Wang, X.; Zhang, J. Effects of straw and biochar amendments on aggregate stability, soil organic carbon, and enzyme activities in the Loess Plateau, China. *Environ. Sci. Pollut. Res.* 2017, 24, 10108–10120.
- Zhang, H., Henderson-Sellers, A., & McGuffie, K. (1996). Impacts of tropical deforestation. Part I: Process analysis of local climatic change. *Journal of Climate*, 9(7), 1497-1517.
- Zhu, K., Ran, H., Wang, F., Ye, X., Niu, L., Schulin, R., & Wang, G. (2022). Conservation tillage facilitated soil carbon sequestration through diversified carbon conversions. *Agriculture, Ecosystems & Environment*, 337, 108080.

REVIEW ARTICLE

Assessments on the Impacts of Climate Change on Food Production, Nutrition, Quality, and Resource Use Efficiency: A Review

Alemu Andualem¹, Tamirat Wato^{1*}, Tariku Goa¹, Natnale Sitotaw², Gutema Urgi³

¹Department of Plant Science, College of Agriculture and Natural Resource, Bonga University, Ethiopia, Bonga Ethiopia

²Department of Mechanical Engineering, School of Mechanical and Industrial Engineering, Addis Ababa University, Ethiopia

³Department of Resource Utilization and Plant Protection, College of Resource and Environmental Science, China Agricultural University, Beijing China

Corresponding Author: Tamirat Wato: email: tamiratwato1@gmail.com

Received: 04 February, 2023, Accepted: 02 April, 2023, Published: 05 April, 2023

Abstract

Nowadays, climate change is a hot issue all over the world which mainly affects crop production and productivity. Thus, it causes food insecurity all over the globe mainly in Sub-Saharan African countries like Ethiopia. This paper provides a comprehensive overview bonded to the appraisal of climate change impacts on nutrition, quality, and resource use, effectiveness using climate, water, and crop yield models. The studies present that climate change models with advanced spatial resolution can be a way forward for coming climate protrusions. The variability of downfall and the adding temperature was a cause of frequent failure and shortage and had a disastrous impact on the livelihood of the people. Climate change exacerbates the enormous being burden of undernutrition. It affects food and nutrition security and undermines current sweats to reduce hunger and promote nutrition. Undernutrition in turn undermines climate adaptability and the managing strategies of vulnerable populations. Climate change is now a global miracle with growth, poverty, food security, and stability counteraccusations. Because of significant dependence on the agrarian sector for product, employment, and import earnings, Ethiopia is seriously hovered by climate change, which contributes to frequent failure, flooding, and rising average temperatures. The most vulnerable sectors to climate variability and change in the country are husbandry, water, and mortal health. Agricultural sectors are severely affected by Climate change; thus, it causes the production and productivity of animals and plants. To enhance productivity, biotechnology in breeding is therefore essential; nonetheless, optimization is needed for every crop and circumstance. While some newly released crop varieties can increase yield and improve resource use efficiency, others can produce crops on marginal land that are sufficient.

Keywords: Climate, Food security, Nutrition, Resource use efficiency, Temperature

Introduction

Currently, climate change is the most the most factor affecting the environment globally, and also its effects will continue in the coming periods (Temesgen et al., 2014). The climate change is due to the various ways in which it

has destructive effects. Natural variables including volcanic eruptions, changes in the Earth's orbital components, and variations in solar output, as well as human-induced factors, namely the release of greenhouse gases, all contribute to global climate change. Although it is not a new phenomenon, the rate at which it is changing now is unparalleled. The average surface temperature of the Earth increased by 0.6–0.2⁰C in the 20th century as reported by the IGP (Intergovernmental Panel) on Climate Change's 3rd Assessment Report (TAR). According to Parry et al. (2005), this trend is predicted to continue, with temperatures increasing by 1.4 - 5.8⁰C by 2100.

On average, the Ethiopian's annual temperature increased by 1.3⁰C, or 0.28⁰C between 1960 and 2006 in every decade. By the 2060s and 2090s, respectively, it is predicted that the mean annual temperature will rise by 1.1 to 3.1⁰C and 1.5 to 5.1⁰C. As reported by McSweeney et al. (2007) reported that in many models, the expected changes under a single emissions scenario range up to 2.1⁰C. Moreover, climatic projections indicated that as a result of global warming, rainfall unpredictability will rise and extreme flooding and droughts will occur more frequently (World Bank, 2010).

The various reports showed that the most common cause of climate change is the emission of Greenhouse Gases. Similarly, the IPCC (Intergovernmental Panel on Climate Change's) (2007) evidence is now overwhelmingly persuasive that the change in the environment mainly the change in climate resulting due to greenhouse gases (GHGs) is real and that the most vulnerable and disadvantaged people will suffer the most. Moreover, the IPCC (2014a), reported that the global change temperature by 2100 on average may vary from 1.8-4.0⁰C. In the case of plants and animals, about 20 to 30% of the species are forecasted to be under extinction due to the rise in temperature by 1.5-2.5⁰C (FAO, 2010; IPCC, 2014a, b), which will have a significant impact on crop production and maintaining food security in underdeveloped nations as reported by Mekuriaw et al. (2014).

The relationship between climate change and maintaining food security through the production of crops and animals has primarily focused on how it affects agricultural sectors ultimately causing the production of food crops. As an example, Gregory et al. (2002) reported that the wheat and rice crops, showed a reduction in crop length and consequently production of wheat crops as a result of heating and yield decreases of around 5% per ⁰C increase beyond 32⁰C for rice. According to Cline's 2007 projection, agricultural productivity will decline globally by 15.9%, falling 19.7% more sharply in developing nations. Similarly, a simulation of maize output in Latin America and Africa for the year 2055 anticipated a 10% total decrease (Jones and Thornton, 2003, Addisu et al., 2020).

Unambiguous evidence of climate system warming includes rising global average sea levels, higher air and ocean temperatures on average, and resulted in extended snow and melting of ice (IPCC 2007). The seasonal mean temperature has risen in several parts of Ethiopia, according to the IPCC (2014a) study. By 2006, Ethiopia's average annual temperature had risen by 1.1–3.1⁰C, and likewise, McSweeney et al. (2007) explained that there were hotter days and also hot nights on average each year. Animal health and food production are both negatively impacted by this.

Currently, the concerns of crop production and being secure in food will be increased due to the effects of climate change, thus it will make more difficult conditions to produce more production of crops and livestock products to maintain food security in the world. Among these, the most powerful and frequently occurring one is natural disasters. Accordingly, Gregory et al. (2008) and UNFCCC (2009) explained that the consequences of natural disasters are profound for underdeveloped countries that are malnourished, impoverished, and still under food security. This paper's goal was to examine how climate change has affected possibilities for adaptation and mitigation as well as crop production, food security, nutrition, quality, and resource use efficiency.

The Impacts of Climate Change

Impact of Climate Change on Nutrition

The influence of climate change on food security, undernourishment, and agricultural productivity in poor nations is the single greatest health risk, according to World Health Organization research, because so many people are affected (Confalonieri et al., 2007). Global hunger and malnutrition risks are increasing due to climate change, impacting food security, livelihoods, health, water, sanitation, and socioeconomic determinants, affecting food access, maternity care, and sanitation (Easterling et al., 2007).

Women, children, and marginalized populations are among those who are the poorest and most at risk of suffering from anticipated climate change effects (World Food Program 2009). They are highly vulnerable to natural disasters, directly depend on resources that are unstable due to climate change, and have limited capacity to adjust or mitigate its effects. Pastoralists, artisanal fishermen, and smallholder and subsistence farmers will be especially exposed to the intricate, regional consequences of climate change (Easterling et al., 2007).

According to the IPCC (2007), there will be 200–600 million more hungry people by 2080, and 24 million more people will be undernourished by 2050 as a result of developing nations' reduced access to calories. Furthermore, it has been projected that in 2050, there will be a relative increase in mild stunting of 1% to 29% due to climate change as opposed to a world without it. Climate change is expected to cause rates of severe stunting to increase by 23% in central sub-Saharan Africa and 62% in South Asia (Lloyd et al., 2011).

Climate change causes direct and indirect effects. It can cause direct effects on the production and productivity of crops and livestock, the food systems of the country, and its food security, whereas, the reduction and varying nutritional values of the products are the indirect effects of it. Many plant crops used by humans have lower protein concentrations as a result of increased carbon dioxide. The amount of atmospheric carbon dioxide that is expected in the next (22nd) century will have a significant impact on plant physiology and growth, which is anticipated to have an impact on agricultural output and food quality. Under warmer and drier conditions, raised CO₂ is anticipated to have a stronger effect on the levels of grain protein (WHO, 2010 and 2013).

Impacts of Climate Change on Health

Scientists have conjectured about the potential effects of climate change on human health due to the correlation between weather-related conditions and seasonality. Two recent studies are White and Hertz-Picciotto (1985) and Haile (1988). There is insufficient scientific evidence to conclusively demonstrate a link between human health and climate change. Human health is impacted by the climate both directly and indirectly. Heat stress, heart problems, preterm delivery, lung ailments including asthma and bronchitis, and infections spread by mosquitoes and ticks are examples of direct consequences. Premature birth, lung disorders associated with smog, and illnesses like pneumonia and influenza are examples of indirect consequences. For instance, Asthma, hay fever, pneumonia, influenza, and other illnesses are associated with particular climates and weather patterns, including winter, which impact the distribution and life cycles of fungi and plants.

Human health may be impacted by climate-related changes in crop and animal production, surface and subsurface water, coastal resources, social and economic conditions, and more. Poorer diets could ensue from decreased food production, and decline of irrigation or agricultural drainage due to increasing the levels of the sea and altered patterns of precipitation could have serious negative effects on human health and the economy (Harrington et al. 1989). The ranges in topographical and vulnerability of humans to numerous factors like diseases may change as a result of increased human migration from one place to another. Human health will typically be negatively impacted by any situation that lowers standards of life (Chappie and Lave 1982).

Many problems still need to be rectified. Without reliable predictive information on the local temperatures, relative humidity, and precipitation amounts, it is impossible to anticipate the health effects. Confounding variables have an impact on human health, some much more so than the weather. Due to the complicated interrelationships between these components (both with weather and among themselves), global climate change may frequently have an impact on two or more factors at once. We lack the knowledge necessary to fully evaluate all the synergistic and compensating effects. Finally, we lack knowledge of the potential social and economic effects that variations in mortality and morbidity brought on by the climate or ozone might produce. We also don't know a lot about the social and financial consequences associated with such effects. It is difficult to find information, in particular, on the productivity losses and out-of-pocket medical expenses linked to rises in morbidity.

Implications of Climate Change in Food Security

The literature has a wealth of information about the negative effects of climate change on East Africa's agricultural industry. Climate change affects agriculture and food production in several ways. By influencing the rise and distribution of incomes, it indirectly impacts food production by influencing the demand for agricultural products (Gregory et al. 2008). Altering agroecological circumstances (e.g., variations in rainfall causing drought or flooding, or variations in temperature causing changes in the length of the growing season), directly impacts food production. In semi-arid and dry regions, the amplification of high temperatures and little precipitation will be the most noticeable effect of climate change on smallholder and subsistence farmers (Mendelson and Dinar, 2010). The fifth report from the IPCC warns that, particularly for farmers in semi-arid regions, climate change in East Africa could worsen food insecurity, cause people to lose their rural livelihoods, and lower agricultural production. The 2013 IPCC report noted that extreme weather events can be dangerous to critical infrastructure networks and services such as emergency response, water and power supply, and healthcare.

The yields of major cereal crops in the African region are expected to be considerably negatively impacted by climate change (Niang et al. 2014). Mild warming rates of 1 to 20⁰C are putting rare ecological systems in jeopardy and may have an impact on water supply, human health, and food production in some regions. According to "worst-case" projections, warming by 2⁰C by the middle of the century might result in losses of 27–32% for maize, sorghum, millet, and peanuts (Schlenker and Lobell, 2010). According to the IPCC, global warming of 4⁰C or more will raise the possibility of severe, all-encompassing, and permanent effects to which it will be challenging to adapt. Because of numerous factors, including land degradation or nutrient deficiency, quick growth of population, and unavailability of adequate technologies such as newly released crop species, plant nourishments, mechanization, and irrigation have sparked the development of agricultural sectors all over the world. These factors are in common in Ethiopia which makes it a great problem for the governments and other development organizations to maintain food security and alleviate the poverty (Mekuriaw et al. 2014). According to Gebreegziabher et al. (2016); Tadesse and Alemayehu (2019) and Tamirat (2019) reported that the agricultural sector plays a vital role by creating job opportunities for the people, as a result, in Ethiopia more than 85% of the populations are highly engaged in these activities, and also, it aids to GDF of the country approximately \$40 billion; it earns 88% of export revenues, and fulfills 73% of the domestic industries depends on agricultural raw materials needs of the country. Therefore, the primary sector that contributes to food security is agriculture because it is a significant means of food and also, it is important in producing excess capital to hasten the social and economic growth of the nation. However, due to unpredictable and irregular rainfall, this sector is highly at risk in the degraded areas and semi-arid areas of the country. Moreover, Zenebe et al. (2011), reported that climate change has a detrimental effect on financial sectors through reducing revenues by worsening agricultural activities. As a result, if this trend continues in the world mainly in developing countries, there will be a decline in salaries by 2050. According to the World Economic

Forum in 2023, By 2050, unchecked climate change might force over 200 million people to migrate, resulting in poverty and undoing decades of development gains. According to the no-total factor productivity-growth scenario model, income is lost due to climate change by about 30% when compared to the no-climate-change baseline.

Impact of Climate Change on Quality and Resource Use Efficiency

Climate change impacts on water resources

The hydrologic cycle is expected to quicken due to global warming, increasing precipitation and evaporation by 7 to 15% on a worldwide scale (Bolin et al. 1986). For many locations, climate models cannot agree on the direction of yearly precipitation change, hence the consequences on local water supplies are uncertain (Fredenck and Gleick 1989). In areas like northern California, where winter snowfall dominates precipitation and spring snowmelt dominates runoff, warmer temperatures may result in more winter rain, earlier spring melting, and seasonal runoff patterns (Gleick 1987a,b).

Water's relative values for alternate uses are likely to change. Variations in the seasonal and yearly availability of water can affect how water is used and reservoir capacity used for irrigation, fish habitat, flood control, and power generation. According to Frederick and Gleick (1989), hydroelectric power may become more appealing as a way to reduce the greenhouse effect while also coping with potential increases in energy needs. Water must be available at current or suitable new locations to produce more hydroelectric power, but water is running out in many parts of the world.

The building of dams, interbasin water transfers, desalination, waste recycling, and weather modification are examples of climate change adaptations that necessitate the development of non-conventional water sources. To increase performance, water managers might spend money on research and technology developments as well as infrastructure improvement plans, but they might not be able to justify these expenditures until climate change plays a major role (Frederick and Kneese, 1989). The possibility of future climate change could encourage further investment in these fields.

Impacts of climate change on Forests, Unmanaged Ecosystems, and Biodiversity

Within a few decades, the anticipated global warming might occur, possibly surpassing the millennium-scale natural rates of forest migration (Batie and Shugart, 1989). If this is the case, stressed-out existing woodlands become more susceptible to disease, pest infestation, and eventually fire (Clark, 1988). They will eventually replace the current forests with new forms of vegetation or forests that have a combination of species (Sedjo and Solomon 1989; Tamirat and Mekides, 2020).

In high latitudes, a lack of summer warmth and a lack of water limit tree development, while in the middle latitudes, heat and a lack of water limit growth. The consequences on forests would likely be negligible in the tropics, where temperature increases are predicted to be the least severe. As a result of climate change, the boreal forests will most likely migrate northward onto the tundra that is not now covered in trees, if there is enough precipitation and suitable soils. The biggest transitions, according to simulations, happen along the boundary between the boreal and cool temperate regions. Especially if higher CO₂ and better plant water use efficiency do not result in the predicted improvements in tree growth and moisture-saving benefits (Tamirat and Mekides, 2020), some mid-latitude forests may perish. According to Sedjo and Solomon (1989), species in mountainous areas would migrate to higher altitudes as temperatures rose.

Ecosystem biodiversity is at risk from rapid climate change, according to Batie and Shugart (1989). Certain extant plant and animal species would not be able to adapt because they are not mobile enough to migrate at the rate required for existence (Davis, 1989a, b). Although it is difficult to quantify, biodiversity has a significant economic worth. The forest industry must remove early species, try to salvage, thin, seed expensively, and actively plant trees in harvested stands to adapt to changing climates (Sedjo and Solomon, 1989).

In comparison to agriculture, the introduction of new types occurs much more slowly in forestry. Changes in the species mix may result from adaptation, at least in the early decades, and may necessitate expensive modifications to the logging and processing industries. Long tree growth cycles further increase the financial risk of selecting the wrong species for the changing climate, discouraging investment in trees and mills to process them. Production forestry will shift geographically, with some areas becoming more and more significant providers of wood products while others experience a loss. Only those locations where high-yield plantation forestry can still be carried out profitably will actively manage their woods.

Other unmanaged terrestrial and freshwater ecosystems have nonmarket value to humans because of their rarity (they might be protected in national parks, for example), significance in maintaining genetic and biotic diversity (Peters and Darling 1985; Graham 1988; Wilson 1988), and general ecological context they provide for natural resources that humans exploit. Research has indicated that the distribution of biotic communities and vegetative life zones, such as grasslands and tundra (Emanuel et al. 1985), arid communities (Neilson, 1986, 1987), and forests (Pickett and White 1985, Overpeck et al. 1990, Tamirat and Mekides, 2020), may be affected by global warming. Mainly the arid regions are particularly vulnerable (Adam et al. 1978; Dregne, 1893). Changes in former climates have been found to have a significant impact on vegetal patterns in pale ecological studies (Davis and Botkin 1985; Webb, 1986; Woodward 1987; Davis, 1989a, b). Concern is growing over how global warming may affect arctic and alpine communities, highly specialized terrestrial species, and species with weak dispersion systems (Peters and Darling 1985). Since aquatic communities are intimately linked to their terrestrial environments through energy, nutrients, and water, changes in terrestrial vegetation could have a substantial impact on freshwater systems, even though the effects of global warming on aquatic communities are still unknown (Minshall et al. 1985; Tamirat, 2019).

Climate Change Adaptation and Mitigation Measures

According to FAO (2010), the most important strategy to maintain food security and reduce the impacts on the environment is Biotechnology. In the meanwhile, modified crop types that can withstand extreme weather conditions including drought, waterlogging, salt, and climate change might increase the area that can be planted with crops, like in eroded soils, to improve the availability of foods for the future.

There is a great deal of worry that the rising levels of greenhouse gases in particular, carbon dioxide contribute to global warming by absorbing long-wave radiation reflected off the earth's surface. Carbon in the atmosphere has increased by 30% over the previous 150 years. According to Stavins and Richards (2005), the majority of scientists concur that elevated atmospheric carbon dioxide concentrations and increasing global temperatures are causally linked.

Increasing the worldwide storage of carbon in soils is one strategy suggested for lowering atmospheric carbon dioxide. However, storing carbon in soil benefits everyone. According to Kumar et al. (2009), Adesodun and Odejim (2010), and others, it boosts soil quality, improves agronomic production, advances food security, and mitigates climate change by offsetting anthropogenic emissions. Programs for conservation and reforestation have been implemented in this scenario throughout the past three decades (Tamirat and Mekides, 2020). To adapt to climate change, smallholder farmers must manage agricultural risk through climate-smart agriculture, enhance

climate information services, and accelerate adaptation over decadal time scales using integrated technology, agronomy, and policy alternatives.

Conclusion

Numerous studies have established the validity of climate change, the likelihood that it will worsen, and the likelihood that the most vulnerable and disadvantaged people will suffer the most. The security of food and nutrition is directly impacted by climate change, undermining present initiatives to combat food insecurity, the vitally important yet less addressed social, economic, and human health-related issues in the world. Human health may be impacted by climate-related changes in agricultural production mainly crop production, aquaculture, water and coastal resources, social and economic conditions, and more. The variability of climate such as the presence of unpredictable rainfall, floods, and droughts; and the variation in temperature, and precipitation can cause an impact on agricultural sectors. To achieve the necessary scale and rate of climate change, the following points should be considered; a) the integration of climate change policies and their implementation, b) the policies and implementations should be evidence-based, c) to maintain food security at all levels of the nations, the usage of climate-smart approach should be mandatory. The major effects of climate change on the yield of crops and livestock feeds, the availability of water, the occurrence of pandemics and unexpected diseases, and flood damage will result from variations in rainfall and rise in temperature. Improved rotation systems, reduced tillage carbon-sequestration practices, and higher crop cover including agroforestry, are just a few of the CSA strategies for climate change adaptation and mitigation that should be strengthened. Construction of additional dams and reservoirs, inter-basin water transfers, and the creation of "unconventional" sources of water, such as desalination, reutilizing of various waste materials from the industry, municipal sectors, and agricultural sectors, as well as weather modification, could all be considered adaptations to climate change.

Acknowledgment: None

Funding: None

Conflict of Interest: All authors declared that no compete of interest exists.

Authors contribution: All authors equally contributed from the concept development up to the final approval of the manuscript.

Data availability: All necessary data were incorporated within the paper.

References

- Adams R, Adams M, Wdlen A. (1978). Drylands: Man and Plants. The Architectural Press Ltd, London.
- Adesodun J, Odejim O. (2010). Carbon-Nitrogen sequestration potentials and structural stability of a Tropical Alfisol as influenced by Pig-Composted Manure. *Journal of International Agrophysics*, 24, 333-338.
- Adisu M, Ayele T, Zerhun G, Ashenafi H. (2020). Climate change impacts on household food security and adaptation strategies in southern Ethiopia. *Food and Energy Security*, 10, 1-14. <https://doi.org/10.1002/fes3.266>.

- Batie SS, Shugart HH. (1989). The biological consequences of climate changes: an ecological and economic assessment. In: Rosenberg, N. J., Easterling, Wm.111, Crosson. P. R., Darmstadter, J. (eds.). Greenhouse Warming: Abatement and Adaptation. Resources for the Future, Washington, D.C., Pp. 121-131.
- Bolin B, Doos BR, Jaeger J, Warrick RA. (eds.) (1986). The greenhouse effect, climatic change, and ecosystems. Scope 29. John Wiley, Chichester. Pp. 541.
- Chappie M, Lave L. (1982). The health effects of air pollution: a reanalysis. *J. Urban Econom*, 12(3), 346-376
- Cline WR (2007). Global warming and agriculture: impacts estimate by country. Centre for Global Development and Peterson Institute for International Economics, Washington, DC.
- Confalonieri U, Menne B, Akhtar R, Ebi KL, Hauengue M, Kovats RS, Revich B, Woodward AJ. (2007). Human health. In: Parry ML, Canziani OF, Palutikof JP, van der Linden PJ, Hanson CE, eds. Climate change, impacts, adaptation and vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge and New York: Cambridge University Press. Pp. 391–431.
- Davis MB. (1989a). Lags in vegetative response to greenhouse warming. *Climate Change*, 15, 75-82.
- Davis MB. (1989b). Insights from paleoecology on global climate. *Bull. Ecol. Soc. Am.*, 70, 222-228.
- Davis MB, Botkin DB. (1985). Sensitivity of cool-temperate forests and their fossil pollen record to rapid temperature change. *Quaternary Res.*, 23: 327-340. [https://doi.org/10.1016/0033-5894\(85\)90039-0](https://doi.org/10.1016/0033-5894(85)90039-0)
- FAO. (2010). World agriculture: towards 2015/2030 summary report. Food and Agriculture Organization, Rome; Financing for food security, adaptation and mitigation. Rome: Food and Agriculture Organization. "Climate-smart" agriculture. Policies, practices and Food and Agroforestry in Cold Deserts of Ladakh, India. Food, Agriculture, and the Environment. The Potential for Soil Carbon Sequestrate.
- Frederick KD, Gleick PH. (1989). Water resources and climate change. In: Rosenberg, N. J., Easterling, Wm. 111, Crosson, P. R., Darmstadter, J. (eds.) Greenhouse warming: abatement and adaptation, Chap. 10. Resources for the Future, Washington, D.C., Pp. 133-143.
- Frederick KD, Kneese AV. (1989). Western water allocation institutions and climate change. In: Waggoner, P. E. (ed.) Climate and water. John Wiley, New York. Pp. 395-41.
- Gebreegziabher Z, Stage J, Mekonnen A, Alemu A. (2016). Climate change and the Ethiopian economy. A CGE analysis. *Environment and Development Economics*, 21(2), 205-225. [doi:10.1017/S1355770X15000170](https://doi.org/10.1017/S1355770X15000170)
- Gleick PH. (1983a). The development and testing of a water balance model for climate impact assessment: modeling the Sacramento Basin. *Water. Resour. Res.*, 23, 1049-1061.
- Gleick PH. (1987b). Regional consequences of increases in atmospheric CO₂ and other trace gases. *Clim. Change*, 10:137-161.
- Gregory PJ, Ingram JSI, Andreson R et al. (2002). Environmental consequences of alternative practices for intensifying crop production. *Agric Ecosyst Environ.*, 88, 279–290.
- Gregory PJ, Ingram JSI, Brklacich M. (2008). The impacts of climate change on food security in Africa. *International Agrophysics*, 24, 333-338.
- Haile DG. (1988). Computer simulation of the effects of changes in weather patterns on vector-borne disease transmission. Report prepared for US EPA, Office of Policy, Planning, and Evaluation. US EPA Project No DW12932662-01-1.
- Harrington W, Irupnick AJ, Spofford WO, Jr. (1989). The economic losses of a waterborne disease outbreak. *J. of Urban Economics*, 25(1), 116-137.
- IPCC. (2013). Climate-smart agriculture sourcebook. Executive summary (Food and Agriculture Organization of the United Nations).

- IPCC (2014a). Climate change; impacts, adaptation, and vulnerability. Part A: summary for policymakers: Global and Sectorial Aspects. In: Field CB et al. (ed) Cambridge University Press, Cambridge.
- IPCC. (2014b). Climate change: mitigation of climate change. In: Edenhofer O et al. (eds) 29, note 4, Cambridge University Press, Cambridge.
- IPCC. (2007). Summary for policymakers. In: Solomon S, Qin D, Manning M, et al., editors. Climate change the physical science basis. Contribution of working group I to the fourth assessment report of the intergovernmental panel on climate change. Cambridge: Cambridge University Press; Pp. 1–18.
- Jones PG, Thornton PK. (2003). The potential impacts of climate change on maize production in Africa and Latin America in 2055. *Global Environ Change*, 13, 51–59.
- Kumar P, Murkute A., Gupta, S, Singh S. (2009). Carbon Sequestration with Special Reference to Badege B. (2001). Deforestation and land degradation in the Ethiopian highlands: A strategy for physical recovery. *Northeast African Studies*, 8(1), 7–26.
- Likens GE. (1985). The aquatic ecosystem and air-landwater interactions. In: Likens, G. E. (ed.) An ecosystem approach to aquatic ecology, Mirror Lake and its environment. Springer-Verlag, New. Pp. 1985 – 516.
- Lipper L, Thornton P, Campbell, B. et al. (2014). Climate-smart agriculture for food security. *Nature Clim Change*, 4, 1068–1072. <https://doi.org/10.1038/nclimate2437>.
- Lloyd SJ, Kovats S, Chalabi Z. (2011). Climate change, crop yields, and under nutrition: development of a model to quantify the impact of climate scenarios on child under nutrition. *Environ. Health Prospect*, 119, 1817–23.
- McSweeney C, New M, Lixcano G. (2007). Ethiopia – UNDP Climate change country profiles. Pp. 27.
- Mekuriaw S, Tegegn F, Mengistu A. (2014). A review on reduction of greenhouse gas emission from ruminants through nutritional strategies. *Acada J Environ Sci.*, 2(1), 006–014.
- Mendelson R, Dinar A. (2009). Climate change and agriculture. An economic analysis of global impact, adaptation and distribution effects. Elgar, Cheltenham. *European Review of Agricultural Economics*, 37(3), 421–423. <https://doi.org/10.1093/erae/jbq027>.
- Minshall GW, Robert C. Petersen Jr, Curtis FN. (1985). Species Richness in Streams of Different Size from the Same Drainage Basin. *The American Naturalist*, 125(1), 16–38. <https://www.jstor.org/stable/2461606>.
- Neilson RP. (1986). High resolution climatic analysis and Southwest biogeography. *Science*, 232, 27–34.
- Neilson RP. (1987). Biotic regionalization and climatic controls in western North America. *Vegetation*, 70, 135–147.
- Niang I, Ruppel OC, Abdrabo MA, Essel A, Lennard C, Padgham J, Urquhart P. (2014). Climate change in Africa: impacts, adaptation and vulnerability. Contribution of working group II to the fifth assessment report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge.
- Overpeck JT, Rnd D, Goldberg R. (1990). Climatic-induced changes in forest disturbance and vegetation. *Nature. Lond.*, 343, 51–53.
- Parry J, Hammill A and Drexhage J (2005). Climate Change and Adaptation. International Institute for Sustainable Development (IISD).
- Peters RL, Darllng DJS. (1985). The greenhouse effect and nature reserves. *BioSci.*, 35(11), 707–717.
- Raloff J. (1989). Global smog: newest greenhouse projection. *Science News*, 135(17), 262–263.
- Schlenker W, Lobell DB. (2010). Robust negative impacts of climate change on African agriculture. *Environ Res Lett.*, 5(1), 014010. IOP Publishing Ltd Stability of a Tropical Alfisol as Influenced by Pig-Composted Manure.

- Sedjo RA, Solomon AM. (1989). Climate and forests. In: Rosenberg, N. J., Easterling, WM. 111, Crosson, P. R., Darmstadter, J. (eds) Greenhouse warming: abatement and adaptation. Resources for the Future, Washington, D.C., Pp.105-119.
- Stavins R, Richards K (2005). The cost of U.S. forest-based carbon sequestration. Prepared for the Pew Center on Global Climate Change, USA. Pp. 52.
- Tadesse A, Alemayehu M. (2019). Impacts of Climate Change on Food Security in Ethiopia: Adaptation and Mitigation Options: A Review. In: Castro, P., Azul, A., Leal Filho, W., Azeiteiro, U. (eds) Climate Change-Resilient Agriculture and Agroforestry. Climate Change Management. Springer, Cham. https://doi.org/10.1007/978-3-319-75004-0_23
- Tamirat W. (2019). Improvements of Crop Production through Integrated Soil Fertility Management in Ethiopia. *Asian Journal of Environment & Ecology*, 11(1), 1-11. DOI: 10.9734/AJEE/2019/v11i130130.
- Tamirat W, Mekides A. (2020). Opportunities and Challenges of Scaling up Agroforestry Practices in Sub-Saharan Africa: A Review. *Agricultural Reviews*, 41(3), 216-226. 10.18805/ag.R-154.
- Temesgen G, Wondie M, Daniel H, Abeba N. (2014). Climate Change Adaptation and Mitigation Measures in Ethiopia. *Journal of Biology, Agriculture and Healthcare*, 4(15), 148-152.
- UNFCCC. (2009). Climate change, food insecurity and hunger. Technical Paper of the IASC Task Force on Climate Change.
- Webb T. (1986). Is vegetation in equilibrium with climate? How to interpret late-Quaternary pollen data. *Vegetation*, 67, 75-91.
- White PS, Plckett STA. (eds.) (1985). The ecology of natural disturbance and patch dynamics. Academic Press, New York.
- Woodward FI. (1987). Climate and plant distribution. Cambridge University Press. Cambridge. Pp. 188.
- World Bank (2010). Economics of Adaptation to Climate Change, Ethiopia. World Bank Group 1818 H Street, NW Washington, DC, Pp. 20433.
- Zenebe T, Kannan S, Yilma D, Beyene G. (2011). Invasive Bacterial Pathogens and their Antibiotic Susceptibility Patterns in Jimma University Specialized Hospital, Jimma, Southwest Ethiopia. *Ethiopian Journal of Health Sci.*, 21(1), 1-8. DOI: 10.4314/ejhs.v21i1.69038.

RESEARCH ARTICLE

Urinary outputs of nickel in association with their concentration levels in water, soil, and selected foods among farmers in the industrial estate of district swabi

Ijaz Ahmad¹, Niamat Ullah^{1*}, Zia-ud-Din¹

¹Department of Human Nutrition, The University of Agriculture Peshawar Pakistan

Corresponding Author: Niamat Ullah, niamatullah@aup.edu.pk

Received: 11 January, 2023, Accepted: 12 March, 2023, Published: 05 April, 2023

Abstract

Diet is the main route of exposure to trace metals, so the assessment risk of these elements to human via dietary intake is important. The non-carcinogenic health risk of Nickel (Ni) to the farmers via dietary intake in the Gadoon Amazai Industrial Estate (GAIE) Swabi of Kyber Pakhtunkwa was assessed. A cross sectional study was carried out in the GAIE to estimate the concentration of Nickel in all types of vegetables, grains, drinking water, irrigation water, soil and in urine of the farmers. A total of 22 farmers, living within the 2km distance in all four directions were selected and enrolled in the study after signing consent form. Atomic absorption spectrophotometer was used for the Ni analysis in the collected samples of water (both drinking and irrigation water), soil, foods and in urine of farmers. Results shows that the mean age of the farmers using tube well water and wastewater for irrigation purposes was 43.5 ± 21.01 and 44.75 ± 16.44 years, height was 167.6 ± 3.7 and 165.75 ± 6.02 cm, weight was 61 ± 12.52 and 64.75 ± 9.63 kg and BMI was 21.65 ± 3.73 and 23.7 ± 4.4 respectively. The concentration of the Ni in the wastewater irrigated field was significantly higher than the tube well irrigated field. The mean concentration of Ni in the soil irrigated with wastewater was 123.50 ± 54.74 mg/kg respectively and in the tube-well irrigated field was 54.25 ± 10.14 mg/kg. The Ni concentration in the wastewater irrigated garlic, fodder grass, potato, wheat and maize were 9.15 ± 0.50 mg/kg, 8.82 ± 1.30 mg/kg, 7.70 ± 1.04 mg/kg and 7.56 ± 1.24 mg/kg respectively compared to tube-well irrigated land i.e., 0.97 ± 0.25 mg/kg, 0.64 ± 0.42 mg/kg, 1.08 ± 0.35 mg/kg, 1.05 ± 0.013 mg/kg and 1.02 ± 0.39 mg/kg. A positive correlation was observed between the water, soil, and all crops grown in the GAIE. The bio-accumulation factor was higher for Ni in both the site. The Hazzard Quotient (HQ) for Ni exceeded the 1 for crops irrigated with industrial wastewater compared with tube-well irrigated crops and thus pose adverse health affect to farmers health. This study concluded that a strong association was evident between the Ni concentration in crops and cereals and waste-water irrigation.

Keywords: Dietary intake; Health risk; Hazzard Quotient (HQ); Farmer's health; Nickel; Industrial Wastewater

Introduction

Heavy metals are found on the earth crust naturally and their exposure to the environment occurs through both human and natural activities. Metals have crucial biological effects to both plants and animals but they became toxic after exceeding a certain limits, when enter to body these heavy metals combine with biomolecule of body (such as proteins and enzymes) form stable bio-toxic compounds (Mahurpawar, 2015).

Heavy metals are not degradable in nature and possess to accumulate in different parts, due to no proper mechanism to eliminate from the body these heavy metals even at low concentration damage the health of both humans and animals (Arora, Kiran, Rani, Rani, Kaur, & Mittal, 2008). Each metals possess specific toxicity signs and their effect may be acute, chronic, or sub-chronic (Hossain, Ahmed, Abdullah, Akbor, & Ahsan, 2015).

Diet is essential component for human body, fruits and vegetables provide nutrients (CHO, Protein, minerals and vitamins) to body, therefore contamination of fruits and vegetables by metals cannot be underestimated (Itanna, 2002). The heavy metals enter to food chain by the consumption of vegetables (Wang, Shan, Zhang, & Wen, 2004). Consumption of unsafe contaminated heavy metals for long time through foodstuff results deposition of metals in the kidney and liver, which causing disturbance in various processes and leads to some kind of diseases like nervous, cardiovascular, kidney and bone (Järup, 2003). Some nutrients are depleted by the intake of contaminated food which causes intrauterine growth retardation, lower immunological defenses, impaired psycho-social behaviour and the gastrointestinal cancer (Arora, Kiran, Rani, Rani, Kaur, & Mittal, 2008). The nature of effect can be neurotoxic, mutagenic, carcinogenic, or teratogenic (Duruibe, Ogwuegbu, & Ekwurugwu, 2007).

In the economy of any nation good health and productive agriculture is important especially in the poverty. Pakistan is an agriculture country, and the farmers are considered to be the backbone of Pakistan economy. Agriculture system can be affected by health of producer's (farmers). The poor health of farmers decreases the work capacity and ability to explore various farming practices and also result in loss of workdays, decrease innovation ability. The agriculture production process and its output effected by both good and poor health of the farmers as well as in the society (Corinna & Ruel, 2006)

Due to the adverse health effects, heavy metals became an important concern in the agricultural products. Heavy metals even at very low concentrations are significantly very toxic due to its cumulative nature. The heavy metals accumulated at toxic level in the crop by the long term application of wastewater to irrigated field (Juste & Solda). Enormous volumes of wastewater is released by rapid urbanization and industrialization, which is used as a source for irrigation practices. The wastewater irrigation are creating problems and opportunities for agriculture production, as this wastewater contains considerable amount of toxic heavy metals and plants nutrients respectively (Singh, Mohan, Sinha, & Dalwani, 2004). The wastewater treatment does not remove the heavy metals and thus causes risk to food chain by heavy metals contamination by soil (Fytianos, Katsianis, Triantafyllou, & Zachariadis, 2001).

Vegetables are grown on small scale as compared with commercial main crops like, maize, wheat, and rice, but the productivity of vegetable totally depends on the good quality water availability for irrigation purposes. However, because of recently increase in the exportation of vegetables to other countries, in Pakistan the area of vegetables cultivation is increasing by the time. Like, vegetables were cultivated during 2007 and 2008 at about 253,800ha (M. Abbas, Parveen, Iqbal, Riazuddin, Iqbal, Ahmed, et al., 2010). However, the most commonly consumed vegetables are grown in the areas of peri-urban where farmers using polluted waste-water for irrigation purpose coming from sewage with no proper filtration. Thus, it is expected that from the peri-urban locations of have polluted irrigation water, all the grown vegetables accumulate a considerable amount of heavy metals (Firdaus-e & Tahira, 2011). A variety of heavy metals are present in wastewater, by extensive uses of contaminated waste-water as source of irrigation for cereals and vegetables accumulate toxic metals (Adhikari, Manna, Singh, & Wanjari, 2004). These heavy metals easily enter the food chain, because the removal of these toxic metals from the water and soil is very difficult (Wilson & Pyatt, 2007).

Several approaches have been proposed to estimate the potential health risks of contaminants, distinguished mainly by carcinogenic and non-carcinogenic effects. United States Environmental Protection Agency (USEPA) current methods of assessing non-cancer risk and cancer risks are very different. The standard cancer risk assessment methods can be used to quantify the magnitude of risk, while similar methods are not available for

quantifying the non-cancer risks (U. EPA, 1989). The non-cancer risk assessment is based on the use of hazards quotient (HQ), which is a ratio of the estimated dose of a contaminant to the Reference Dose (reference dose or RfD is the level below which there will not be any appreciable risk of the contaminant). If the estimated dose for an exposed population is equal to or greater than the RfD, then the population is at risk of contracting the adverse effect associated with the contaminant (U. EPA, 1989; USEPA). To assess the overall potential risk of more than one contaminant for non-carcinogenic effects, a Hazard Index (HI) approach has been developed based on the United States Environmental Protection Agency (Urban & Cook, 1986) Guidelines for Health Risk Assessment of Chemical Mixtures. The HI represent the total non-cancer hazard for all exposure pathways presented. The HI is equal to the sum of all the hazard quotients in USEPA-1989 (A. EPA, 1989). When the hazard index exceeds unity, there may be concern for potential health effects. Any single contaminants with exposure level greater than toxicity value will cause the hazard index to exceed unity, the hazard index can also be exceeded for multiple chemicals even if no single chemical exceed its RfD (A. EPA, 1989). We have carried out this study to estimate the concentration of Nickel (Ni), and to assess the hazard quotient and target hazard quotient of the Ni via the foodstuff for the local farmers in the Industrial Estate of District Swabi. A variety of heavy metals are present in wastewater, by extensive uses of contaminated wastewater as source of irrigation for cereals and vegetables accumulate toxic metals (Adhikari, Manna, Singh, & Wanjari, 2004). These heavy metals easily enter the food chain because the removal of these toxic metals from the water and soil is very difficult (Wilson & Pyatt, 2007). The target hazard quotients (THQs) and hazard index (HI) were calculated to assess non-carcinogenic health effects from individual and combined heavy metals because of daily foodstuff consumption.

Material and Methods

Study site

The Gadoon Amazai Industrial Estate (GAIE) is in district Swabi, Khyber Pakhtunkhwa, and is 325 meters elevated above the sea level, bounded north, east, west and south by Baisak, Maini, Gandaf, and Topi. GAIE contains a total of 330 active units and was established in 1986-1987 (Khan, Ahmad, Shah, Rehman, & Khaliq, 2009). The main active units in the industrial state are steel, marble, textiles, chemicals, soap and soap, plastic, and ghee and cooking oil (Khan, Ahmad, Shah, Rehman, & Khaliq, 2009).

Inclusion Criteria and sample size

All those farmers having farms within 2 kilometers from industrial estate were orientated about the study. A total of 22 farmers were randomly selected and data were collected from the agreed farmers.

Anthropometric Assessment:

Farmer's body weight and height were assessed through WHO standard methods. Weight was measured through digital scale, while weighing, the farmers were asked to remove heavy clothes, shoes and all un-necessary things. The height of farmers were measured through studio-meter. BMI was calculated through the height and weight data.

Sampling and pre-treatment

The study area was divided into 4 directions (North, South, West and East) from the mid-point. The samples of water, soil, crops, vegetables (Potato, Garlic) Milk, and Urine were selected for the study. Water of both the irrigation and drinking purpose were collected, for drinking purpose water from different sources (Tube-well, hand-pump, Open-well and tape water) were used in the Industrial estate. The farmers use two types of water for irrigation of fields 1) Industrial wastewater and 2) Tube-well water. The water samples and milk of animals were directly collected in polythene bottles, all the bottles were washed with acidify water and dried, the morning urine samples were collected with the addition of 2 drops of HCL to reduce the decomposition of urine by bringing its PH to below 4. The crops samples which were grown in the industrial area were collected in polythene bags, labelled, and brought in ice-cold boxes to the Department of Human Nutrition, The University of Agriculture Peshawar for further analysis. In the laboratory the vegetable samples were thoroughly washed firstly by ordinary tape water followed by distilled water to eliminate soil and air burn pollutant. Edible portion of vegetables samples were cut into pieces and placed in oven for 48 hours at 80⁰ C to completely dry. After the heating the samples were kept overnight to cool down to room temperature. Each sample was ground to fine powder and stored for further chemical treatment.

Digestion and treatment

All the required glassware was first washed with standard detergent followed by tap water, soaked in an acid bath (10% HCL) and placed in oven to dry. 1 gm of sample was taken into the digestive tube, 12 ml concentrated nitric acid was added to the tubes and kept overnight. The next day 4 ml of perchloric acid was added and placed in heat-block, gradually increased the temperature from 80⁰ C until a white fume started AOAC, 2000 (Cuniff, 2003). After heating the solution was cooled down at room temperature filtered in 50 ml volumetric flask, diluted by distilled water up to the mark (50 ml). The powdered soil was treated with solutions of HNO₃, perchloric acid and sulphuric acid with a 4:1:1, samples were filtered using Whatman No.42 filter paper to eliminate suspended substances. Prepared samples were stored in the clean bottle prior analysis (Shakya & Khwaounjoo, 2013). Atomic absorption spectrophotometer was used as standard method of AOAC 2000(Cuniff, 2003) for the determination of nickel in the samples.

Health Risk Assessment:

The health risk from the consumption of contaminated food was calculated by the following quotient hazards equation (Huang, Zhou, Sun, & Zhao, 2008) (Muhammad, Ullah, & Jadoon, 2019).

$$HQ = \frac{CDI}{RfDo}$$

$$CDI = \frac{CF \times IR \times EF \times ED}{BW \times AT}$$

Where CDI is the chronic daily intake of metal from food expressed in mg kg⁻¹ day⁻¹, RfDo is the oral reference dose (in mg kg⁻¹ day⁻¹). CF is the concentration of metal in plants, IR is the ingestion rate, EF is the exposure frequency (365 days per year), ED is the exposure duration, BW is the body weight and AT is the averaging

exposure time(364×ED). The RfDo values for Ni was 0.02 mg/kg according to US-EPA. If the HQ values exceed the unity, there will be potential effect to the body.

Daily Intake of Metal:

The daily intake of metals (DIM) from the food sources was calculated by the following equation (Bi, Zhou, Chen, Jia, & Bao, 2018) (Orisakwe, Nduka, Amadi, Dike, & Bede, 2012).

$$DIM = \frac{C_{fveg} \times W_{veg}}{BW}$$

Where C_{veg} is the concentration of metal in in vegetables or food (mg/kg), W_{veg} (mg/day) is the ingestion rate of food contaminated by particular metal and BW is the Body weight (kg)

Bio-accumulation Factor:

The bio-accumulation factor also known as transfer factor, which is the transfer of heavy metals from the soil to the crops grown in an area. The bio accumulation factors is an index reflecting the ability of a plant species to accumulate a particular metal regards to its concentration in the soil (Galal & Shehata, 2015; Ghosh & Singh, 2005) It is calculated as the metal concentration in the plants (dried weight basis) divided by the concentration of that particular metal in the soil on which it grows (Cui, Zhu, Zhai, Chen, Huang, Qiu, et al., 2004; Liu, Zhao, Ouyang, Söderlund, & Liu, 2005), (Ahmad, Khan, Ashfaq, Ashraf, & Yasmin, 2014), (Muhammad, Ullah, & Jadoon, 2019).

$$TF = \frac{CP}{CS}$$

Where CP is the concentration of metal in plants and CS is the concentration of metal in the soil. The bio-accumulation shows the bio-availability of metals and nutrients to the plants.

Health Risk Assessment:

The health risk from the consumption of contaminated food was calculated by the following quotient hazards equation (Huang, Zhou, Sun, & Zhao, 2008) (Muhammad, Ullah, & Jadoon, 2019).

$$HQ = \frac{CDI}{RfDo}$$

$$CDI = \frac{CF \times IR \times EF \times ED}{BW \times AT}$$

Where CDI is the chronic daily intake of metal from food expressed in $\text{mg kg}^{-1} \text{ day}^{-1}$, RfDo is the oral reference dose (in $\text{mg kg}^{-1} \text{ day}^{-1}$). CF is the concentration of metal in plants, IR is the ingestion rate, EF is the exposure frequency (365 days per year), ED is the exposure duration, BW is the body weight and AT is the averaging

exposure time($364 \times ED$). The RfDo values for the Ni is 0.02 mg/kg according to US-EPA. If the HQ values exceed the unity, there will be potential effect to the body.

Results and discussion

The body mass index (BMI) gives the best gauge for estimate of the nutritional status of the farmers, As the farmers characterize an occupational group which required heavy amount of physical activity for their field work in farm. Because of this heavy workload the farmers might tend to decrease risk for development of overnutrition. The farmers use tube well water for irrigation purposes had BMI within normal range ($18.5\text{--}24.9 \text{ kg/m}^2$) as compared to farmers uses wastewater for irrigation as shown in table 1. Different studies reported different obesity status of farmers in different countries like in Australia only 15.2% of the farmers reported to overweight (Dorner, Leitner, Stadlmann, Fischer, Neidhart, Lawrence, et al., 2004) and in Greece 86.1% of the farmers reported to overweight (Vardavas, Linardakis, Hatzis, Saris, & Kafatos, 2009).

Table 1. Anthropometric, Socio-demographic features of the farmers

Variables		Tube Well Mean $\pm$ SD/Frequency (%)	Wastewater Mean $\pm$ SD/Frequency (%)
Age (Years)		43.50 $\pm$ 21.01	44.75 $\pm$ 16.44
Weight (kg)		61 $\pm$ 12.5	64.75 $\pm$ 9.63
Height (cm)		167.5 $\pm$ 3.9	165.8 $\pm$ 6.02
BMI (kg/m^2)		21.61 $\pm$ 3.73	23.7 $\pm$ 4.4
Household size		8.25 $\pm$ 1.3	7.25 $\pm$ 1.5
Area of Farming (kanals)		62.5 $\pm$ 28.7	42.00 $\pm$ 33.9
Living since (Years)		8.25 $\pm$ 6.0	11.25 $\pm$ 6.1
Educational Level	Illiterate	70%	50%
	Literate	30%	50%

The farmers living in the study location were not of permanent resident that's why the mean time of residency for both the categories of farmers were 8.25 ± 6.0 and 11.25 ± 6.1 years. Education is positively related to the production of farmers the literacy rate of farmers in the study location was very low. Because of poverty and workload in the field, the farmers are unable to enroll their children for education. The literacy rate of farmers in underdeveloped and developing countries remain the lowest as reported in Odisha (Das & Sahoo, 2012). In south Nigeria about 33.8% of the farmers were reported uneducated (Fabunmi, Aba, & Odunaiya, 2005).

The industries effluents were fallen directly to nearest small canal of water, theses small canals were then used for the irrigation purposes, so the waste from the industries were transferred through canals water to the farming land. The study area was divided in to four directions (North, East, West and South) from the mid-point to estimate the Ni in all the industries effluents in irrigation water.

Figure 1 shows the Nickel concentration in the irrigation water sources in the GAIE. The wastewater used for irrigation purposes had significantly high concentration of nickel compared to the tube-well irrigated water. The highest mean of 3.87 ppm nickel was present in the North side followed by East 3.57 ppm, West 2.19 ppm and South 1.87 ppm in the wastewater sources for irrigation. No nickel was found in the East site in tube-well irrigation source. The nickel concentration in the waste-water irrigation crossed the permissible limit of 0.2 ppm

set by the US Irrigation Water Quality standards, while in the tube-well irrigation water were in the safe limit for irrigation.




Figure 1. Nickel concentration in the irrigated water

This high amount of nickel was due to the nickel-cadmium batteries, steel and ghee and oil, kitchen appliances, surgical instruments and steel alloys industries in GAIE (Tariq, Ali, & Shah, 2006). These industries released a vast amount of nickel to the environment. Industrial activities such as mining, electroplating, and manufacturing of essential commodities produce a huge volume of wastewater as effluents containing heavy metals and other toxicants, which deteriorate the quality of aquatic system (S. Abbas, Sarfraz, Mehdi, & Hassan, 2007), (Bose & Bhattacharyya, 2008). In the study area, drinking water was consumed from the four types of sources, from tube-well, tape water hand pump and well-water in the home.




Figure 2. Nickel concentration in the Drinking water

Figure 2 shows the concentration of nickel in the drinking water sources used in the Industrial zone of GAIE. The results showed the Ni concentrations was significantly higher in the open well-water used for drinking. The Ni concentration in tube-well and hand pump drinking sources was found within the safe limit of 0.05 ppm set by the Pakistan National Standard for Drinking water (NSDWQ-Pak). The highest concentration of Ni contamination was found in the Well-water in West and South site of 0.1 ppm, while 0.09 ppm and 0.07 ppm in North and East site. No nickel was detected in the tube-water in the East, South and North regions (detection limit for nickel is 0.02). The highest concentration in the tap-water was recorded in the North and South region of 0.06 ppm followed by West and East of 0.05 ppm and 0.05 ppm respectively. Compared with concentration of Ni of 0.037 ppm in Karachi (Karim, 2011) the current concentration was higher.

Table 2 illustrate the nickel concentration in the soil, wheat, maize, potato, garlic and fodder grass grown in the industrial estate by the waste-water irrigation. The mean concentration of nickel in all the vegetables, soil and milk samples in waste-water irrigation was significantly higher than in the tube-well irrigation. The nickel concentration in soil samples irrigated by tube-well and wastewater was measured as 54.25 ± 10.14 mg/kg and 123.50 ± 54.74 mg/kg. In the tube-well irrigated wheat the Ni concentration was in range from 0.93 to 1.2 mg with a mean of 1.05 ± 0.13 mg compared to the wastewater irrigated wheat of range from 6.14 mg to 8.93 mg. Whereas the mean concentration of Ni in the maize was recorded as 1.02 ± 0.39 mg in the tube-well irrigated compared with the wastewater irrigated of 6.31 ± 0.76 mg.

Table 2. Concentration of Nickel in Food, grown in the industrial estate irrigated through Tube Well and Industries-wastewater

Variables	Tube Well Source	Wastewater Source	p-value
Soil	54.25 ± 10.14	123.50 ± 54.74	.047
Wheat	1.05 ± 0.13	7.56 ± 1.24	.001*
Maize	1.02 ± 0.39	6.31 ± 0.76	.004*
Potato	1.08 ± 0.35	7.70 ± 1.04	.002*
Garlic	0.97 ± 0.25	9.15 ± 0.50	.003*
Fodder	0.64 ± 0.42	8.82 ± 1.30	0.00**

*Significant difference observed at $p < 0.05$ ** Significant difference observed at $p < 0.01$

Among all the food sources in the waste-water irrigation, the highest mean value of Ni was recorded in the garlic 9.15 ± 0.50 mg, while in the tube-well sources was recorded in the potato samples of 1.08 ± 0.35 mg. The Ni mean concentration in the wastewater irrigated potato was 7.70 ± 1.04 mg while in the tube-well irrigated was 1.08 ± 0.35 mg. The average value in the fodder grass was recorded as 8.82 ± 1.30 mg in the water-water irrigation compared to tube-well irrigation of 0.64 ± 0.42 mg.

The results of this study is resembles with the finding of (Hussain, Khattak, Shah, & Ali, 2015) who studies the contamination of soils by the industrial effluents, and resulted that the mean concentration of nickel in the waste-water soil was 119.8mg. compared with the reference or tube-well irrigated soil which was 58.8mg. The highest concentration in the industrial waste-water was due to the stainless steel, alloys industries, which direct expelled their waste to the environment without any treatment. A study (Al-Othman, Ali, Al-Othman, Ali, & Habila, 2016) reported that the mean nickel concentration in the wheat grains grown by tube-well water in district Swabi was 0.087mg lower than the finding of this study. The average concentration of Ni in the wheat and maize irrigated with un-polluted water was 0.04 ± 0.01 mg and 0.11 ± 0.01 mg, significantly lower than that of the crops grown with

the polluted water which was 0.1 ± 0.01 mg in wheat and 0.10 ± 0.01 mg in maize. The soil contamination with toxic metals and pathogens was due to the long term irrigation with waste-water (Farahat & Linderholm, 2013). The extractable concentration of nickel in the present study was lower than the soil irrigated by canal water and waste-water of Hyderabad city in southern Pakistan (Jamali, Kazi, Arain, Afridi, Jalbani, & Memon, 2007). Nickel is utilized in certain industrial applications and can be a potential contaminant in food products. Ni is also an essential element for human health but may become toxic above certain levels. There is currently no published permissible limit of Ni in milk; however, researchers still maintain a focus on Ni contamination in milk due to its potential for negative health impacts (Ismail, Riaz, Akhtar, Goodwill, & Sun, 2019) and the concentration of nickel in milk was found in the range of 0.0070-2.631 mg/l.




Figure 3. Nickel Concentration in Urine

Figure 3 shows the Ni concentration in the urine of farmers using food from wastewater irrigated land and tube-well irrigated land. The waste-water farmers urine showed significantly high concentration of nickel compared to the farmers urine using tube-well water for irrigation. The highest concentration of 1.5 ppm was observed in the farmers urine of West side to industries, while the lowest was recorded in the North side farmers, who were using wastewater for the irrigation purposes. While in the farmers who was using tube-well water for irrigation, the highest mean concentration 0.81 ppm of nickel was recorded in the farmers urine to the East side to industries, followed by the West, South and North. According to the Korea National survey for environmental pollutants in the human body 2018, and heavy metals in the blood and urine of the Korean population (Lee, Lee, Moon, Choi, Lee, Yi, et al., 2012) reported that the high level of heavy metals in the urine and blood was due to the high intake of contaminated foods, the level of toxic metals in urine was strongly related to its oral intake.

The bio-accumulation and transfer factor is used to estimate the plants potential to attract the particular metal or nutrients from the soil through roots (Farahat & Linderholm, 2015) (Galal & Farahat, 2015). Table 4.3 shows the

bio-concentrations factors of Ni for crop-soil system. The crops grown on Wastewater possess higher accumulation factors compared with crops grown on tube-well water.

Table 3. Nickel content for tube well and wastewater for different crops

Crops	Nickel	
	Tube well	Wastewater
Wheat	0.021±0.006	0.050±0.014
Maize	0.021±0.012	0.042±0.012
Potato	0.022±0.011	0.051±0.013
Garlic	0.020±0.008	0.033±0.011
Fodder grass	0.014±0.011	0.057±0.10

Table 4. Correlation between Soil and Crops

Soil crops	Ni
Soil-Wheat	.807*
Soil-maize	.771*
Soil-potato	.813*
Soil-garlic	.725*
Soil-Fodder	.802*
Fodder-Milk	.987**

* Correlation is significant at the 0.05 * Correlation is significant at the 0.01

Table 4 shows the Pearson correlation among the soil and crops, soil and fodder and fodder to milk was determined to check the route of the nickel from soil to crop, fodder and from fodder to milk. It was found that a strong correlation (significant at 0.05) was found for nickel.

Health risk is defined as the quotient between the estimated to daily metal intake from the soil through food chain and oral reference dose for each metal. An index under the value 1 assumed as safe. The hazard quotient from nickel is presented in figure 4.

The HQ value for nickel exceed the unity in the food grown by waste-water irrigation water and pose adverse health effect to farmers health as shown in figure 4.4. All the foods grow with tube-well water pose no health effect from nickel.

The present study resemble with the study (Qin, Zou, & Qiu, 2008) stated that the health risk of chromium was less compared to nickel because of the high RfDo value of 1.5mg/day. They calculated the health risk of the heavy metals to the general public of Guangzhou China. There result showed the HQ value for chromium in the safe limit of below the unity.




Figure 4 . Hazard Quotient for Nickel

Conclusion

The wastewater from the industries in the Gadoon Amazai Industrial Estate (GAIE) contained significant amount of nickel. Soil, vegetables, cereals and fodder for cattle from the agriculture land irrigated through wastewater from GAIE contained significant amount of nickel. Food consumption had been identified as the major pathway of human exposure to heavy metals. Farmers of GAIE area shows significant amount of nickel in their Urine. The health risk assessment indicates that the farmers of GAIE were at high risk of nickel toxicity.

Declaration

Acknowledgment: The authors would like to acknowledge the support of Pakistan Atomic Energy Commission for their support in laboratory analysis.

Funding: N/A

Conflict of interest: N/A

Authors contribution: N/A

Data availability: From the authors

References

- Abbas, M., Parveen, Z., Iqbal, M., Riazuddin, M., Iqbal, S., Ahmed, M., & Bhutto, R. (2010). Monitoring of toxic metals (cadmium, lead, arsenic and mercury) in vegetables of Sindh, Pakistan. *Kathmandu University Journal of Science, Engineering and Technology*, 6(2), 60-65.
- Abbas, S., Sarfraz, M., Mehdi, S., & Hassan, G. (2007). Trace elements accumulation in soil and rice plants irrigated with the contaminated water. *Soil and Tillage Research*, 94(2), 503-509.

- Adhikari, T., Manna, M., Singh, M., & Wanjari, R. (2004). Bioremediation measure to minimize heavy metals accumulation in soils and crops irrigated with city effluent. *Journal of Food Agriculture and Environment*, 2, 266-270.
- Ahmad, K., Khan, Z. I., Ashfaq, A., Ashraf, M., & Yasmin, S. (2014). Assessment of heavy metal and metalloid levels in spinach (*Spinacia oleracea* L.) grown in wastewater irrigated agricultural soil of Sargodha, Pakistan. *Pak J Bot*, 46(5), 1805-1810.
- Al-Othman, Z. A., Ali, R., Al-Othman, A. M., Ali, J., & Habila, M. A. (2016). Assessment of toxic metals in wheat crops grown on selected soils, irrigated by different water sources. *Arabian Journal of Chemistry*, 9, S1555-S1562.
- Arora, M., Kiran, B., Rani, S., Rani, A., Kaur, B., & Mittal, N. (2008). Heavy metal accumulation in vegetables irrigated with water from different sources. *Food chemistry*, 111(4), 811-815.
- Bi, C., Zhou, Y., Chen, Z., Jia, J., & Bao, X. (2018). Heavy metals and lead isotopes in soils, road dust and leafy vegetables and health risks via vegetable consumption in the industrial areas of Shanghai, China. *Science of the Total Environment*, 619, 1349-1357.
- Bose, S., & Bhattacharyya, A. (2008). Heavy metal accumulation in wheat plant grown in soil amended with industrial sludge. *Chemosphere*, 70(7), 1264-1272.
- Corinna, H., & Ruel, M. (2006). Understanding the links between agriculture and health. *Published by International Food Policy Research Institute, Brief*, 1.
- Cui, Y.-J., Zhu, Y.-G., Zhai, R.-H., Chen, D.-Y., Huang, Y.-Z., Qiu, Y., & Liang, J.-Z. (2004). Transfer of metals from soil to vegetables in an area near a smelter in Nanning, China. *Environment International*, 30(6), 785-791.
- Cuniff, P. (2003). Official Methods of Analysis of AOAC International, 17th edn, 2nd revision. *Gaithersburg, MD, USA: AOAC International*.
- Das, A. B., & Sahoo, D. (2012). Farmers' educational level and agriculture productivity: a study of tribals of KBK districts of Odisha. *International Journal of Education Economics and Development*, 3(4), 363-374.
- Dorner, T., Leitner, B., Stadlmann, H., Fischer, W., Neidhart, B., Lawrence, K., Kiefer, I., Rathmanner, T., Kunze, M., & Rieder, A. (2004). Prevalence of overweight and obesity in Austrian male and female farmers. *Sozial-und Präventivmedizin/Social and Preventive Medicine*, 49(4), 243-246.
- Duruibe, J. O., Ogwuegbu, M., & Ekwurugwu, J. (2007). Heavy metal pollution and human biotoxic effects. *International Journal of physical sciences*, 2(5), 112-118.
- EPA, A. (1989). Risk assessment guidance for superfund. Volume I: human health evaluation manual (part a). In): EPA/540/1-89/002.
- EPA, U. (1989). Risk assessment guidance for superfund. Vol. 1: Human health evaluation manual. EPA/se0/1-89/002.
- Fabunmi, A., Aba, S., & Odunaiya, N. (2005). Prevalence of low back pain among peasant farmers in a rural community in South West Nigeria. *African journal of medicine and medical sciences*, 34(3), 259-262.
- Farahat, E., & Linderholm, H. W. (2013). Effects of treated wastewater irrigation on size-structure, biochemical products and mineral content of native medicinal shrubs. *Ecological engineering*, 60, 235-241.
- Farahat, E., & Linderholm, H. W. (2015). The effect of long-term wastewater irrigation on accumulation and transfer of heavy metals in *Cupressus sempervirens* leaves and adjacent soils. *Science of the Total Environment*, 512, 1-7.
- Firdaus-e, B., & Tahira, S. A. (2011). Metal accumulation potential of wild plants in tannery effluent contaminated soil of Kasur, Pakistan: field trials for toxic metal cleanup using *Suaeda fruticosa*. *J Hazard Mater*, 186(1), 443-450.
- Fytianos, K., Katsianis, G., Triantafyllou, P., & Zachariadis, G. (2001). Accumulation of heavy metals in vegetables grown in an industrial area in relation to soil. *Bulletin of environmental contamination and toxicology*, 67(3), 0423-0430.

- Galal, T. M., & Farahat, E. A. (2015). The invasive macrophyte *Pistia stratiotes* L. as a bioindicator for water pollution in Lake Mariut, Egypt. *Environmental monitoring and assessment*, 187(11), 701.
- Galal, T. M., & Shehata, H. S. (2015). Bioaccumulation and translocation of heavy metals by *Plantago major* L. grown in contaminated soils under the effect of traffic pollution. *Ecological Indicators*, 48, 244-251.
- Ghosh, M., & Singh, S. (2005). A comparative study of cadmium phytoextraction by accumulator and weed species. *Environmental Pollution*, 133(2), 365-371.
- Hossain, M. S., Ahmed, F., Abdullah, A. T. M., Akbor, M. A., & Ahsan, M. A. (2015). Public health risk assessment of heavy metal uptake by vegetables grown at a waste-water-irrigated site in dhaka, bangladesh. *Journal of Health and Pollution*, 5(9), 78-85.
- Huang, M., Zhou, S., Sun, B., & Zhao, Q. (2008). Heavy metals in wheat grain: assessment of potential health risk for inhabitants in Kunshan, China. *Science of the Total Environment*, 405(1-3), 54-61.
- Hussain, R., Khattak, S. A., Shah, M. T., & Ali, L. (2015). Multistatistical approaches for environmental geochemical assessment of pollutants in soils of Gadoon Amazai Industrial Estate, Pakistan. *Journal of Soils and Sediments*, 15(5), 1119-1129.
- Ismail, A., Riaz, M., Akhtar, S., Goodwill, J. E., & Sun, J. (2019). Heavy metals in milk: global prevalence and health risk assessment. *Toxin Reviews*, 38(1), 1-12.
- Itanna, F. (2002). Metals in leafy vegetables grown in Addis Ababa and toxicological implications. *Ethiopian Journal of Health Development*, 16(3), 295-302.
- Jamali, M., Kazi, T., Arain, M., Afridi, H., Jalbani, N., & Memon, A. (2007). Heavy metal contents of vegetables grown in soil, irrigated with mixtures of wastewater and sewage sludge in Pakistan, using ultrasonic-assisted pseudo-digestion. *Journal of Agronomy and Crop Science*, 193(3), 218-228.
- Järup, L. (2003). Hazards of heavy metal contamination. *British medical bulletin*, 68(1), 167-182.
- Juste, C., & Solda, P. Heavy Metal Availability in Long-term Experiments. *Factors influencing sludge utilization practices in Europe*, 8-10.
- Karim, Z. (2011). Risk assessment of dissolved trace metals in drinking water of Karachi, Pakistan. *Bulletin of environmental contamination and toxicology*, 86(6), 676-678.
- Khan, S., Ahmad, I., Shah, M. T., Rehman, S., & Khaliq, A. (2009). Use of constructed wetland for the removal of heavy metals from industrial wastewater. *Journal of environmental management*, 90(11), 3451-3457.
- Lee, J. W., Lee, C. K., Moon, C. S., Choi, I. J., Lee, K. J., Yi, S.-M., Jang, B.-K., Jun Yoon, B., Kim, D. S., & Peak, D. (2012). Korea National Survey for Environmental Pollutants in the Human Body 2008: heavy metals in the blood or urine of the Korean population. *International journal of hygiene and environmental health*, 215(4), 449-457.
- Liu, W.-h., Zhao, J.-z., Ouyang, Z.-y., Söderlund, L., & Liu, G.-h. (2005). Impacts of sewage irrigation on heavy metal distribution and contamination in Beijing, China. *Environment International*, 31(6), 805-812.
- Mahurpawar, M. (2015). Effects of heavy metals on human health. *Int. J. Res. Granthaalayah*, 1-7.
- Muhammad, S., Ullah, R., & Jadoon, I. A. (2019). Heavy metals contamination in soil and food and their evaluation for risk assessment in the Zhob and Loralai valleys, Baluchistan province, Pakistan. *Microchemical Journal*, 103971.
- Orisakwe, O. E., Nduka, J. K., Amadi, C. N., Dike, D. O., & Bede, O. (2012). Heavy metals health risk assessment for population via consumption of food crops and fruits in Owerri, South Eastern, Nigeria. *Chemistry Central Journal*, 6(1), 77.
- Qin, W.-s., Zou, X., & Qiu, R. (2008). Health risk of heavy metals to the general public in Guangzhou, China via consumption of vegetables. *Journal of Agro-Environment Science*, 27(4), p1638-1642.
- Shakya, P. R., & Khwaounjoo, N. M. (2013). Heavy metal contamination in green leafy vegetables collected from different market sites of Kathmandu and their associated health risks. *Scientific World*, 11(11), 37-42.
- Singh, K. P., Mohan, D., Sinha, S., & Dalwani, R. (2004). Impact assessment of treated/untreated wastewater toxicants discharged by sewage treatment plants on health, agricultural, and environmental quality in the wastewater disposal area. *Chemosphere*, 55(2), 227-255.

- Tariq, M., Ali, M., & Shah, Z. (2006). Characteristics of industrial effluents and their possible impacts on quality of underground water. *Soil Environ*, 25(1), 64-69.
- Urban, D., & Cook, N. (1986). Standard evaluation procedure for ecological risk assessment. Washington DC: USEPA. In): EPA-540/9-85-001.
- USEPA, U. Environmental Protection Agency. 2000. Stressor Identification Guidance Document. In): EPA 822-B-00-025. Office of Water and Office of Research and Development
- Vardavas, C. I., Linardakis, M. K., Hatzis, C. M., Saris, W. H., & Kafatos, A. G. (2009). Prevalence of obesity and physical inactivity among farmers from Crete (Greece), four decades after the Seven Countries Study. *Nutrition, Metabolism and Cardiovascular Diseases*, 19(3), 156-162.
- Wang, X.-p., Shan, X.-q., Zhang, S.-z., & Wen, B. (2004). A model for evaluation of the phytoavailability of trace elements to vegetables under the field conditions. *Chemosphere*, 55(6), 811-822.
- Wilson, B., & Pyatt, F. B. (2007). Heavy metal bioaccumulation by the important food plant, *Olea europaea* L., in an ancient metalliferous polluted area of Cyprus. *Bull Environ Contam Toxicol*, 78(5), 390-394.

REVIEW ARTICLE

A review of agroforestry as a sustainable and resilient agriculture

Asif Raihan*

Institute of Climate Change, Universiti Kebangsaan Malaysia, Bangi 43600, Selangor, Malaysia

Corresponding author: Asif Raihan, Email: asifraihan666@gmail.com

Received: 28 March, 2023, Accepted: 22 May, 2023, Published: 26 May, 2023

Abstract

The agricultural sector is confronted with the formidable challenge of providing sustenance for a global population of 9 billion individuals by the year 2050, all the while mitigating adverse ecological and societal impacts. An attempt to address this difficulty has been made through the implementation of organic farming practices, which have yielded predominantly favorable results. Nevertheless, there are still certain obstacles that need to be addressed. Organic agricultural practices exhibit lower yields compared to conventional methods, while concerns persist regarding greenhouse gas emissions and fertilizer leaching. This paper provides an overview of existing organic and conventional agriculture systems and proposes that agroforestry, a deliberate integration of trees and shrubs with crops or livestock, may represent a promising avenue for advancing sustainable agriculture. Agroforestry possesses the capacity to sustain productivity and concurrently provide many ecosystem services through the use of nature-inspired methods. This study presents an overview of the prevalent methods and products associated with agroforestry, while also highlighting the positive environmental and social impacts it brings about. The present study aims to examine the obstacles encountered in the implementation of agroforestry practices and to suggest potential strategies for policy modification that could enhance the uptake of such practices among farmers. The findings of this review study indicate that agroforestry emerges as a very effective land use strategy for addressing both food security and environmental degradation concerns.

Keywords: Agroforestry; Organic agriculture; Land use; Agroecology; Food security; Environmental sustainability

Introduction

The field of agriculture exerts a significant influence on the Earth's ecosystem (Bishaw et al., 2022; Raihan, 2023a). According to Ahmed and Ambinakudige (2023), around 38% of the Earth's land surface is allocated for agricultural purposes, rendering it the most significant anthropogenic land utilization. The primary driver of deforestation and the subsequent loss of native habitats is the expansion of agricultural land (Begum et al., 2020; Jayathilake et al., 2021). This phenomenon has resulted in the collapse of various wildlife populations, including avian, insect, and mammalian species, some of which are currently classified as endangered (Shah et al., 2022). The process of nutrient leaching from fertilizer contributes to the phenomenon of eutrophication in water bodies, hence causing the formation of oxygen-depleted areas known as "dead zones" in various aquatic environments globally (Zahoor & Mushtaq, 2023). Agriculture stands as the primary anthropogenic source of greenhouse gas emissions, which have been linked to climate change. The impacts of these repercussions are not exclusive to the human population.

Derouiche et al. (2023) have observed the presence of detectable levels of pesticides in several habitats, including the human body. The economic burden of pesticide poisoning in the United States has been approximated to be \$1.2 billion annually (Donley, 2019). Additionally, the presence of excessive nitrate in drinking water resulting from over-fertilization can lead to health issues and necessitate costly remediation efforts (Zahoor & Mushtaq, 2023). In addition to the potential environmental and societal consequences, the resilience of our agricultural systems is also a matter of concern (Dipu et al., 2022; Raihan & Tuspekova, 2022a). According to Xu et al. (2022), a mere fifteen crops are responsible for generating 90% of the global food calorie supply. Among these crops, wheat, rice, and maize alone contribute to 60% of the total food calories. The cultivation of the majority of these crops predominantly occurs in expansive areas characterized by yearly monocultures, hence posing a significant susceptibility to pest and disease outbreaks (Khatri et al., 2023). The Irish potato famine, which occurred from 1845 to 1850, resulted in the loss of more than one million lives. This historical event serves as a poignant illustration of the consequences that might arise when a singular crop, upon which a population substantially depends, is devastated by disease (Read, 2022). Monocultures necessitate annual replanting, substantial resource inputs, and weed management (Zhang et al., 2023). It has been posited that this repetitive pattern of planting, fertilizing, and spraying primarily benefits major agribusiness corporations that provide the necessary inputs, rather than effectively addressing global food security objectives (Gerhardt et al., 2022). In order to ensure the enduring viability of an agricultural system, it is imperative that the soil maintains its productivity and that the essential inputs remain accessible in subsequent periods (Paul et al., 2023). Nevertheless, it is important to note that in numerous agricultural settings, the rate of soil loss surpasses that of soil formation, leading to a deterioration in the quality of the remaining soil (La et al., 2023). The agricultural sector is vulnerable to variations in fuel pricing and supplies due to its significant dependence on fossil fuels, particularly in the form of liquid fuel and fertilizer (Majeed et al., 2023; Raihan, 2023b). Simultaneously, the unidirectional flow of fertilizer nutrients contributes to both pollution and scarcity. Phosphorus serves as a pertinent illustration in this context. This indispensable nutrient for plants is projected to undergo a rise in mining and processing costs. Concurrently, the discharge of phosphorus into water bodies contributes to the phenomenon of eutrophication (Mancho et al., 2023).

In the foreseeable future, it is anticipated that our agricultural systems will need to undergo adjustments in response to a shifting climate, characterized by an augmented occurrence of severe weather phenomena such as droughts and floods (Brick et al., 2022; Raihan, 2024a). Furthermore, there is an expected rise in the prevalence of diseases and pests affecting agricultural production (Balasundram et al., 2023). The impact of climate change is expected to be more pronounced in developing regions due to socioeconomic factors such as poverty, which might impede individuals' capacity to effectively respond and adapt (Bedeke, 2023; Raihan, 2023c). The Dust Bowl phenomenon that occurred during the 1930s serves as an illustrative case of the detrimental outcomes resulting from unsustainable farming methods in conjunction with an unprecedented period of severe aridity (Yuan et al., 2023). The collapse of civilizations, such as the ancient Mesopotamians and the Mayans, can be attributed to agricultural overreach and the failure to effectively respond to climate change. The agricultural sector is confronted with the formidable challenge of providing sustenance to a global population of 9 billion individuals by the year 2050, all the while mitigating detrimental impacts on the environment and society (Wijerathna-Yapa & Pathirana, 2022; Raihan, 2023d). Hence, the primary objective of this study is to critically examine the existing organic and conventional agricultural systems, while proposing agroforestry as a potential advancement towards achieving sustainable agriculture. Agroforestry possesses the capacity to sustain productivity and concurrently provides many ecosystem services through the use of systems that imitate the activities observed in nature. This study provides an overview of the prevalent methods and products associated with agroforestry, while also highlighting the positive environmental and social impacts that result from its implementation. This study additionally examines the obstacles encountered in the implementation of agroforestry practices and investigates potential strategies for

modifying policies to enhance farmer engagement and uptake. This review study could provide valuable insights for the development of agricultural and environmental policies aimed at promoting food security through effective land use management, while also mitigating environmental degradation and the adverse effects of climate change.

Methodology

The objective of this study is to assess the potential impact of agroforestry on global sustainability, with a specific focus on ensuring food security and improving environmental quality. The present study employed the systematic literature review methodology as suggested by Tawfik et al. (2019). The systematic literature review framework is considered to be a dependable approach (Benita, 2021; Raihan & Bijoy, 2023; Raihan & Himu, 2023; Raihan, 2023e; 2023f; 2023g). A preliminary review of the literature was conducted to identify pertinent articles, validate the proposed idea, avoid redundancy with previously covered issues, and ensure the availability of sufficient articles for conducting a comprehensive analysis of the subject matter. Moreover, the focal point of the themes should revolve around the inquiries on the significance of agroforestry in ensuring food security and improving environmental quality. According to Tawfik et al. (2019), it is crucial to enhance the retrieval of results by acquiring a comprehensive understanding and familiarity with the study topic through the examination of pertinent materials and active engagement in relevant debates. This objective can be achieved by conducting a thorough examination of pertinent literature and actively participating in pertinent academic conversations.

The present study examined various strategies aimed at mitigating the influence of prejudice. One of the methods employed was performing a systematic manual search to identify any document that might have been missed during the original search process. This investigation, employing the methodology employed by Vassar et al. (2016), discovered no discernible indications of bias. In the context of this investigation, a comprehensive set of strategies was employed to carry out manual searches. The method employed encompassed many strategies, such as conducting an exhaustive literature search to identify relevant references from the studies and reviews under consideration. Furthermore, supplementary materials, including related papers and articles cited within reputable academic databases such as Google Scholar, Scopus, and Web of Science, were thoroughly examined. The manual search results were initially enhanced and polished through the process of examining the reference lists of the included publications. The initial stage of the process was undertaken. Subsequently, the author engaged in the practice of citation tracking, a method involving the systematic monitoring of all the scholarly works that reference each of the papers incorporated in the collection. In conjunction with the manual search, an online search of databases was also undertaken as an integral component of the comprehensive search process.

This study exclusively relied on research articles published in peer-reviewed journals, ensuring the reliability and validity of the findings. The results of this study serve as a valuable basis for future research endeavors that aim to explore the potential impact of agroforestry on achieving sustainability, with a specific focus on ensuring food security and improving environmental quality. Both qualitative and quantitative secondary literature on agroforestry were considered. The publications were thereafter evaluated to ascertain whether their main subject matter bore a resemblance to that of the present inquiry. Priority consideration was given to papers published after the year 2000. The primary justifications for the elimination of papers are their lack of relevance, duplication, incomplete textual content, or limited presence of abstracts. The predetermined exclusion criteria were established to safeguard the researcher against potential biases that could influence their findings. Figure 1 illustrates the progression of review criteria employed for the selection of suitable documents for review analysis.




Figure 1. The development of criteria for the selection of documents.

The initial search with the keywords led to 4463 documents. After scanning the documents based on the selection criteria and to remove possible duplication, 512 articles were selected for the next step of scanning. After screening those article's titles and abstracts, the comprehensive literature review encompassed a total of 146 distinct scholarly articles. The present study implemented a data verification process, wherein each included article was cross-checked with its corresponding entry in an extract sheet using visual evidence. It is noteworthy to mention that of the 146 papers subjected to qualitative synthesis, only those publications containing relevant material were cited in the reference list contained in the manuscript. This implies that certain articles were not included in the reference list. Figure 2 illustrates the systematic review procedure utilized in the current study. After the research topic was chosen, this study proceeded to find and locate relevant articles, do an analysis and synthesis of diverse literature sources, and create written materials for article review. The synthesis phase encompassed the collection of a wide range of publications, which were subsequently amalgamated into conceptual or empirical analyses that were relevant to the finalized research.




Figure 2. The procedure of systematic review conducted by the study.

Organic Farming and Associated Challenges

The emergence of organic agriculture can be attributed to its role as a viable alternative to the prevailing conventional farming paradigm. While there may be subtle variations across countries and certification agencies, the primary principles governing organic management generally entail the prohibition of synthetic pesticides and fertilizers, genetically modified organisms (GMOs), and the preventive application of antibiotics in cattle feed (Mie et al., 2017). In order to preserve soil quality, it is imperative to employ a range of strategies, including but not limited to crop rotation, cover cropping, and mulching (Crystal-Ornelas et al., 2021). According to Davis et al. (2022), animals that are managed under organic practices are required to consume feed that has been certified as organic. Additionally, ruminant animals must be provided with a designated amount of time to access pasture. The maintenance of fertility in agricultural systems is commonly achieved by the utilization of leguminous cover crops, the application of organic materials like manure and compost, the use of biologically generated inputs such as blood and feather meal, and the incorporation of mined mineral compounds (Tei et al., 2020).

The management of weeds in organic grain and vegetable systems typically involves the utilization of tillage as a primary control method. However, it is worth noting that cover cropping and crop rotation also hold significant importance in disrupting weed cycles (Pantović & Sečanski, 2023). The management of pests involves the implementation of strategies such as the provision of suitable habitats for advantageous predators, the careful selection of plant stock that exhibits resistance, and the utilization of biologically produced pesticides as a final recourse, if necessary (Monteiro & Santos, 2022). The implementation of organic production standards typically results in better sustainable outcomes in practical applications (Raihan et al., 2023a). According to Beaumelle et al. (2023), organic farms have been found to support greater levels of biodiversity compared to conventional farms. This enhanced biodiversity encompasses various organisms such as insects, plants, soil biota, as well as avian and larger animal species. According to Pergner and Lippert (2023), organic farms frequently exhibit greater diversity in their cropping systems as a result of including livestock and implementing longer crop rotations. According to Monteiro and Santos (2022), the implementation of mechanical and cultural control strategies in managing weeds and pests can result in residual populations at reduced levels, hence promoting biodiversity. According to Prout et al. (2021), the implementation of organic management practices has been observed to enhance soil quality, as indicated by measures of soil organic matter. However, it is worth mentioning that several studies have reported the highest levels of soil quality in the context of no-till conventional agriculture (Montgomery et al., 2022). While conventional farming often outperforms organic farming in terms of yields, research has demonstrated that in drought years, the situation can be reversed. This phenomenon is related to the superior water retention capabilities of soils managed under organic practices (Martín-Lammerding et al., 2021). In general, organic production demonstrates a lower energy consumption per production unit as a result of the elevated energy expenditures associated with conventional fertilizers and pesticides (Mousavi et al., 2023).

It is important to acknowledge that while organic certification imposes strict criteria for the application of pesticides, synthetic fertilizers, and GMO technologies, both conventional and organic growers have access to a diverse range of techniques that can yield positive environmental results. Cover cropping, integrated pest management, the application of manure and composts to enhance soil organic matter, crop rotation, and the integration of livestock and crops are crucial strategies that warrant careful consideration when assessing their benefits. Several studies have examined the comparison between organic and conventional crop systems and have found that the observed enhancements in organic management can be attributed to specific practices such as the application of manure and the implementation of cover cropping. These practices, which are integral to the organic system, have the potential to yield similar benefits if adopted in a conventional system (Scavo et al., 2022; Chinthalapudi et al., 2023). Figure 3 illustrates the fundamental concepts and resultant impacts of organic farming.




Figure 3. The fundamental concepts and resultant impacts of organic farming.

Nevertheless, despite the commendable goals underlying organic certification methods, it is worth mentioning that a significant number of organic crop production systems employ similar fundamental methodologies as conventional farming, hence potentially resulting in comparable adverse outcomes (Telwala, 2023). The practice of producing annual monocultures, which necessitates annual replanting, the use of fertilizers, rigorous weed management, and the employment of mechanical equipment, has exhibited limited alteration, particularly when implemented on a broader scale beyond local market gardens (Feng et al., 2022). The substitution of conventional instruments with less detrimental alternatives is evident in the adoption of organic seeds in place of genetically modified organism (GMO) seeds, the utilization of cultivation or mulch as alternatives to pesticides for weed management, and the implementation of cover crops and manure as substitutes for fertilizers derived from fossil fuels (Caporali, 2021). While these modifications have the potential to reduce environmental impacts, it is important to note that complete elimination of such problems may not be achievable.

The phenomenon of nitrogen leaching serves as a pertinent illustration of environmental consequences that are not completely eradicated. While several studies indicate a potential reduction in nitrate leaching when employing organic management practices, it is important to note that the resulting quantities of nitrate may still provide a risk of contributing to groundwater pollution. Pimentel and Burgess (2014) conducted a comparison of three different rotations that varied in terms of nitrogen sources. These rotations included an organic rotation that incorporated legume cover crops, an organic rotation that utilized animal manures, and a conventional rotation that relied on synthetic fertilizers. The researchers discovered that the leachate samples from all three treatments occasionally surpassed the regulation threshold of 10 ppm for nitrate concentration in potable water. According to Pimentel and Burgess (2014), the nitrogen given to the crops in the form of nitrate experienced a reduction of 20% in the organic animal rotation, 32% in the organic legume rotation, and 20% in the conventional rotation. The leaching of nitrogen

from organic sources was found to be higher compared to that of conventional fertilizers. According to Valenzuela (2023), the application of manures and legume cover crops resulted in the highest nutrient release during periods of fallow or during times that did not align with the crop's nitrogen need.

While it is acknowledged that organic management can lead to an enhancement in soil quality compared to conventional management (Lin et al., 2023), it is important to note that the utilization of tillage for weed control and the incorporation of biomass from cover crops still pose hazards of soil loss and degradation (Francaviglia et al., 2023). The deleterious impacts of tillage, as demonstrated by Pearsons et al. (2023), encompass compaction, erosion, and a reduction in soil biological activity. According to the findings presented in the study conducted by Arnhold et al. (2014), research examining erosion rates in organic and conventional agricultural systems has yielded inconsistent outcomes, which can be attributed to factors such as the specific crop rotation, types of crops employed, and the methods of tillage employed. According to Arnhold et al. (2014), the study conducted by the authors in mountainous regions of Korea revealed that soil erosion rates, regardless of whether conventional or organic management practices were employed, were deemed excessive and unsustainable for long-term productivity. There has been a growing interest in the adoption of no-till techniques for organic farming due to the recognition of the advantages associated with minimizing tillage (Szczepanek et al., 2023). The typical procedure involves cultivating a cover crop before the primary cash crop, followed by mechanically crushing it and afterward planting through the resulting residue (Lamichhane et al., 2023). When executed accurately, the application of mulch effectively inhibits the growth of weeds, hence eliminating the necessity for cultivation in relation to the specific crop. Nevertheless, cultivating the requisite biomass in the cover crop to achieve efficient weed management might pose a formidable obstacle, and the feasibility of this approach may be constrained in arid regions where the cover crop competes for scarce water resources (Nosratti et al., 2023). Perennial weeds present a distinct challenge due to their inherent ability to penetrate and thrive within mulch layers (Ruch et al., 2023).

According to Gou et al. (2022), when considering the measurement of organic systems on a per-area basis, they may exhibit superior performance compared to conventional systems. However, it is important to account for the yield gap in organic systems, as this factor may contribute to higher emissions per unit of output. According to Jones et al. (2023), the rise in soil carbon levels in annual systems is accompanied by the emission of other gases, such as nitrous oxide, which counteract the potential benefits by contributing to climate change. The potential environmental ramifications of differences in yields between organic and conventional systems are also of significance. It is widely acknowledged in academic discourse that organic systems tend to exhibit lower productivity, typically resulting in a reduction in yield ranging from approximately 20% to 25%. However, scholarly literature indicates that this range can vary significantly, spanning from 5% to 50%, contingent upon factors such as the specific crop, soil conditions, level of management intensity, and the methodologies employed in the respective studies (Prairie et al., 2023; Santoni et al., 2023). Critics contend that the implementation of organic management practices would necessitate the allocation of additional land for agricultural purposes to ensure the sustenance of global food security. The consequences of this action would include deforestation and the subsequent loss of habitats, resulting in an adverse environmental outcome (Wijerathna-Yapa & Pathirana, 2022; Raihan & Tuspekova, 2022b). In light of the aforementioned issues inherent in the ongoing discourse between organic and conventional approaches, it appears prudent to explore alternative methodologies and strategies that could potentially offer viable resolutions. Rather than adopting a binary perspective when considering our agricultural landscapes, it may be more advantageous to embrace a mindset that incorporates both options, sometimes referred to as a "yes-and" approach. Numerous scholars have advocated for the use of a multidisciplinary and multifunctional framework in the design of agroecosystems, as evidenced by the works of Taylor and Lovell (2021), Thiesen et al. (2022), and Stokes et al. (2023). When considering the challenge of simultaneously providing sustenance to the global population and ensuring the long-term viability of the planet,

Foley aptly asserts that a singular approach is inadequate for addressing all the associated issues. Consider the utilization of silver buckshot rather than relying solely on a silver bullet.

Agroforestry Systems

Agroforestry is a multifunctional method that involves the deliberate integration of trees and shrubs with crops or livestock within our food system. The sustainable agricultural technique of agroforestry has been acknowledged for around 50 years (Aryal et al., 2023). The integration of trees into the agricultural landscape is a concept that has existed since the inception of land cultivation. Agroforestry has been found to yield several advantageous results. These include the mitigation of nutrient and pesticide runoff, the sequestration of carbon, the enhancement of soil quality, the control of erosion, the improvement of wildlife habitat, the reduction of fossil fuel consumption, and the promotion of resilience in the context of an unpredictable agricultural future (Sollen-Norrlin et al., 2020; Temegne et al., 2021; Jinger et al., 2022). In summary, the incorporation of trees and other perennial vegetation into a landscape has the potential to alleviate the adverse impacts associated with agricultural practices. Agroforestry exhibits significant potential as a land use plan in both developed and developing regions due to its capacity to concurrently deliver economic, ecological, and cultural advantages (Telwala, 2023; Viñals et al., 2023). Furthermore, agroforestry has the capacity to provide a wide range of goods including timber, crops, fruits, nuts, mushrooms, forages, cattle, biomass, Christmas trees, and herbal medicine (Sollen-Norrlin et al., 2020). A comprehensive assortment of products in a portfolio would facilitate the distribution of revenue streams over different time horizons. These items encompass short-term options such as crops, pasture, livestock, mushrooms, and certain fruits like currants. Additionally, medium-term possibilities include nuts, fruits such as apples or persimmons, biomass, and medicinal plants. Lastly, long-term prospects involve lumber and the potential for increasing property value. The presence of a wide range of products can potentially mitigate risks for farmers while necessitating innovative marketing strategies (Jacquet et al., 2022; Raihan & Tuspekova, 2022c).

Various forms of agroforestry are implemented in different regions worldwide. The field of tropical agroforestry has historically received greater attention and has been more extensively implemented compared to temperate agroforestry. According to Piato et al. (2021), shade-grown coffee and tea systems have undergone significant advancements, and the presence of manual labor renders some tropical agroforestry approaches more feasible compared to regions where mechanized harvesting is prevalent. Agroforestry has held significant cultural significance in indigenous tropical regions as well as temperate locales such as Europe. However, the prevalence of land abandonment and agricultural intensification in northern territories has resulted in a reduction of conventional agroforestry methods (Nair et al., 2021). There exist five widely acknowledged agroforestry approaches, including alley cropping, silvopasture, riparian buffers, windbreaks, and forest farming (Bishaw et al., 2022). These approaches are applicable across diverse farming systems, topographical features, and climatic regions. Figure 4 illustrates various agroforestry approaches.

Alley cropping

Alley cropping is a sustainable agricultural practice that entails cultivating row crops within the spaces between rows of trees (Gagliardi et al., 2022). Trees have the potential to be cultivated for the purpose of producing lumber or fruits and nuts. On the other hand, alley crops encompass a diverse range of cereals, vegetables, or forages that can be harvested for hay. The cultivation of crops yields immediate financial returns, whereas the growth of trees generates money over an extended period of time.

Types of Agroforestry



Alley Cropping

Planting crops between rows (alleys) of trees or shrubs



Forest Farming

Cultivating crops under the protection of a tree canopy



Silvopasture:

Integrating trees on grazing land for livestock



Windbreaks:

Shielding plants and livestock from wind with trees or shrubs



Riparian Forest Buffers:

Reducing nutrient pollution and stabilizing riverbanks by planting trees or shrubs next to streams and rivers

Graphic by Emilie Austin, EESI



Figure 4. Different types of agroforestry practices.

The potential for enhanced production can arise from the interactions between tree and crop species, facilitated by their distinct ecological niches (Fahad et al., 2022). An illustrative instance can be found in a study conducted in France, which demonstrated the favorable compatibility between walnuts and winter wheat due to their distinct growth periods and divergent rooting depths. According to the findings of Dupraz et al. (2021), it was determined that the integrated cultivation technique yields a 40% higher product output per unit area compared to the separate cultivation of the two crops.

Forest farming

Forest farming encompasses a range of activities, including the cultivation of mushrooms, the collection of medicinal herbs such as ginseng and goldenseal, and the commercialization of woody ornamental resources (Chamberlain et al., 2019). The agroforestry method described in this study typically takes place within mature forests that have been cultivated for lumber production, enabling the generation of income without significant disruption (Frey et al., 2023). The management of forest farming systems can vary in intensity, with the level of management determined by the specific product being cultivated and the target market preferences (Raihan & Tuspekova, 2022d). As an illustration, the cultivation of ginseng in woodland environments necessitates substantial pre-planting measures such as site preparation, application of fertilizers, tillage practices, and the use of fungicides. While these interventions have the potential to enhance crop yields, they also incur higher costs and thus introduce greater financial and operational uncertainties. In contrast, the cultivation of wild-simulated ginseng may encompass the practice of gently displacing fallen foliage, sowing seeds, and allowing the ginseng to mature over a span of multiple years prior to its eventual harvest (Yousefi et al., 2020).

Silvopasture

The practice of silvopasture involves the deliberate integration of cattle within a carefully planned combination of trees and pastureland. Silvopasture distinguishes itself from conventional woodland grazing practices by implementing a deliberate arrangement of trees that ensures adequate sunlight penetration for the underlying fodder vegetation, while simultaneously preventing any detrimental impact on the trees caused by animals. According to Smith et al. (2022), the presence of trees provides animals with shelter by offering shade during the hot summer months and reducing wind exposure during the cold winter season. Furthermore, it has been observed that the quality of pasture in areas with partial shade may exhibit an improvement, but with a minor decrease in biomass productivity (Hidalgo-Galvez et al., 2022). According to Poudel et al. (2022), there is no significant difference in the weight gains of livestock when comparing silvopasture with open pasture grazing systems. According to Huang et al. (2023), if the trees are cultivated for lumber purposes as well, it is expected that the farmer's long-term financial performance will enhance without compromising the current level of production.

Windbreaks

Windbreaks, sometimes referred to as shelterbelts, were promptly acknowledged as a valuable agroforestry technique. Windbreaks play a crucial role in mitigating wind erosion, supporting wildlife habitats, and enhancing water availability for adjacent crops through reduced evapotranspiration and snow capture effects (Subbulakshmi et al., 2023). According to Mallareddy et al. (2023), an increased water supply has the potential to enhance agricultural productivity, hence yielding significant economic advantages for farmers. Windbreaks have the potential to mitigate the heating and cooling requirements of residential and occupational areas on a farmstead by minimizing the infiltration of outdoor air induced by wind (Mume & Workalemahu, 2021). The initiation of the Prairie States Forestry Project by the U.S. government was a response to the Dust Bowl years in North America, aiming to establish a substantial shelterbelt spanning from Canada to Texas (Li, 2021). Another noteworthy illustration pertains to the Three-North Shelter Forest Program in China, which stands as the most extensive afforestation endeavor globally. The initiative commonly referred to as "China's Great Green Wall" was initiated in 1978 and is projected to reach completion by 2050 (Gravesen & Funder, 2022). Comparable approaches have been utilized in Russia, the northern regions of Europe, Australia, New Zealand, and many other nations.

Riparian forest buffers

Riparian buffers refer to vegetated zones established along watercourses that are susceptible to erosion, nutrient leaching, or habitat degradation (Fahad et al., 2022). Typically, there exist two to three distinct "zones" of vegetation, which exhibit variations in their composition as influenced by factors such as proximity to the waterway, slope, and the requirements of primary producers (Lind et al., 2019). Riparian zones have limited suitability for agricultural production, rendering them highly suitable for alternate utilization. The United States Department of Agriculture (USDA) has made a deliberate and coordinated endeavor to enforce conservation practices in the vicinity of water bodies, owing to their advantageous effects on the quality of water and soil. The Environmental Quality Incentive Program (EQIP), administered by the Natural Resources Conservation Service (NRCS), and the Conservation Reserve Program (CRP), managed by the Farm Service Agency (FSA), serve as illustrations of government-funded efforts.

It is important to highlight that, among the five practices mentioned, alley cropping and silvopasture are commonly implemented on land that is deemed appropriate for conventional agriculture. Despite this, it is common practice to engage in conventional cropping for multiple years until the trees reach their full maturity (Dasgupta et al., 2023). Riparian buffers, windbreaks, and forest farming typically manifest in the periphery of fields or on land that is unsuitable for agricultural use. However, it is worth mentioning that in certain instances, the allocation of a portion of cropland may be necessary to achieve the desired width for optimal efficacy (Englund et al., 2021). Hence, these methods have a tendency to serve as a supplement rather than a rival to current production systems, perhaps offering avenues to enhance food security through the utilization of underutilized resources. Agroforestry has the potential to make significant contributions to both conventional and organic agricultural systems in practical applications. In both scenarios, the advantageous impacts of agroforestry have the potential to enhance environmental results beyond the existing capabilities of each respective system. Agroforestry has the potential to mitigate several of the aforementioned issues associated with organic agriculture, such as soil erosion, emissions of greenhouse gases, and leaching of nutrients.

Agroforestry Benefits

The practice of agroforestry has been found to have beneficial impacts on both soil and water quality. The enhancement of soil quality is facilitated by heightened amounts of organic matter, greater diversity in microbial populations, and enhanced nutrient cycling, hence potentially augmenting crop output and bolstering resilience against drought conditions (Fahad et al., 2022). The incorporation of agroforestry vegetative buffer strips has been found to reduce non-point source pollution from row crops, resulting in improvements in water quality (Zahoor & Mushtaq, 2023). Fahad et al. (2022) observed that the implementation of agroforestry and grass buffer strips had a significant impact on reducing the loss of phosphate and nitrogen from a corn-soybean cycle. Perennial vegetation exhibits the capacity to enhance above-ground biomass, thereby impeding runoff and effectively capturing up to 95% of susceptible sediment from being lost (Liu & Lobb, 2021). Additionally, the subterranean roots of these plants have the ability to absorb 80% or more of surplus nutrients, while concurrently serving as a habitat for microbial communities capable of metabolizing pesticides (Behera et al., 2021).

The augmentation of soil organic matter through carbon content not only enhances soil health but also contributes to the mitigation of atmospheric carbon dioxide, which is recognized as a significant factor in climate change (Paul et al., 2023; Raihan et al., 2022a; Raihan & Tuspekova, 2022e; Raihan & Tuspekova, 2023a). According to Lugo-Pérez et al. (2023), the inclusion of trees and shrubs in an agricultural landscape results in a higher degree of carbon sequestration when compared to a monoculture of crops or grassland. In a study conducted by Kim et al. (2016),

a meta-analysis was performed to assess the impact of agroforestry on greenhouse gas emissions. The findings of the study revealed a significant reduction in emissions, with an average mitigation rate of 27 ± 14 tons of CO_2 per hectare per year. Approximately 70% of carbon sequestration was attributed to biomass, while the remaining 30% was sequestered in the soil. According to a study conducted by Udawatta and Jose (2011) in North America, the implementation of agroforestry methods on a small scale has the potential to store around 548.4 Tg of carbon annually. This amount is significant enough to offset almost 34% of the carbon emissions produced by the United States from the burning of coal, oil, and gas. The strategies for enhancing carbon sequestration encompass improved erosion management, heightened carbon storage in woody perennial plants, diminished decomposition of organic matter, and the limited harvesting of crop biomass in agroforestry systems compared to conventional systems (Sivaranjani & Panwar, 2023). The potential significance of the relationship between perennial systems and climate change should not be overlooked. The study conducted by Robertson et al. (2020) examined the possible impact of various annual and perennial systems on global warming. The research findings indicate that all of the annual cropping systems examined, including conventional, no-till, reduced input, and organic, did not result in a reduction of global warming potential. While the farming techniques did indeed result in the accumulation of carbon in the soil, the benefits derived from this were counteracted by the emissions of nitrous oxide. Nevertheless, the implementation of perennial and early successional forest treatments, such as the utilization of alfalfa, hybrid poplar, and the restoration of abandoned early successional sites, resulted in a notable decrease in global warming potential. As the mid-successional and late-successional systems progressed in their development, the annual carbon storage capacity exhibited a decline. The study's findings indicated that the early successional forest system emerged as the most effective strategy for mitigation (Raihan & Tuspekova, 2022f; Raihan & Tuspekova, 2023b). Numerous agroforestry practices demonstrate a high degree of resemblance to the characteristics and dynamics of early successional forests.

The mitigation of climate change is furthered by the adoption of an additional significant strategy, namely the reduction of fossil fuel consumption (Raihan & Tuspekova, 2022g; Raihan et al., 2023b; Raihan, 2023h; Raihan, 2024b). The utilization of bioenergy presents a potential solution for mitigating reliance on fossil fuels (Raihan, 2023i; 2023j). However, apprehensions arise regarding the allocation of important arable land for cultivating energy crops instead of food crops (Kalogiannidis et al., 2023). At present, a significant proportion of the corn yield in the United States, specifically 40%, is allocated towards the production of ethanol. This allocation raises concerns over its potential counterproductivity in relation to the overarching objective of alleviating global food scarcity. According to Ntawuruhunga et al. (2023), the integration of biomass production from trees and food cultivation on the same site holds potential as an agroforestry approach to provide a sustainable energy future without compromising food production capacities. The Land Equivalent Ratio (LER) is a valuable metric for evaluating the comparison between polycultures, which consist of mixed species, and individual crops. This metric takes into account the productivity of the polyculture and computes the land area that would be necessary if the individual crops were cultivated independently. In a study conducted by Haile et al. (2016), a comparison was made between loblolly pine and switchgrass mixes and pure stands of each crop. The researchers observed that while the individual crop yields were lower in the mix, the overall Land Equivalent Ratio (LER) reached 1.47. This implies that cultivating switchgrass and loblolly pine individually would necessitate an additional 47% of land compared to the agroforestry system in order to produce an equivalent quantity of biomass. The Yield-SAFE (Yield Estimator for Long-term Design of Silvoarable Agroforestry in Europe) model was employed to simulate agroforestry systems in Europe. The results of this modeling exercise indicated that integrating trees and crops in Spain, France, and the Netherlands led to higher productivity, as evidenced by the predicted LER (Land Equivalent Ratio) values ranging from 1 to 1.4. This finding was reported by Graves et al. (2007). In a separate study conducted in Switzerland, the implementation of agroforestry models centered around walnut (*Juglans hybrid*) and wild

cherry showed that, in 12 out of 14 instances, the integration of multiple crops resulted in a higher Land Equivalent Ratio (LER) above unity. Furthermore, Sereke et al. (2015) revealed that a significant majority, specifically 68%, of the financial scenarios in Switzerland exhibited higher profitability compared to existing methods.

Agroforestry demonstrates a greater capacity for biodiversity conservation in comparison to conventional and organic monocultures. The inclusion of trees, bushes, and other permanent vegetation within an agricultural landscape has been found to enhance wildlife habitat, resulting in increased abundance and higher diversity of wildlife populations (Ntawuruhunga et al., 2023). In addition to possessing inherent worth, biodiversity has the potential to offer valuable services. According to Ghosh et al. (2023), an increased presence of avian species and predatory insects can effectively regulate pest populations. The provision of suitable habitats for pollinator species has been found to positively impact the pollination of horticulture crops, as demonstrated by Miñarro et al. (2023). According to Ribas et al. (2023), there is typically a decline in the occurrence of diseases in populations that exhibit higher levels of diversity, encompassing both plant and wildlife species. Agroforestry can also provide advantages for livestock. According to Smith et al. (2022), windbreaks serve as a protective barrier against strong winds, safeguarding animals from their adverse effects. Additionally, the provision of shade by trees can enhance thermal comfort during the summer season, potentially promoting more uniform grazing patterns across a paddock. According to Kumar et al. (2023), the implementation of forest-based foraging systems for poultry and hogs has the potential to reduce reliance on grain and create environments that closely resemble the natural habitats of these animals. The cork oak dehesas found in the Mediterranean region exemplify a multifunctional landscape that has persisted for several centuries. These landscapes serve as a source of sustenance for grazing cattle through the provision of grass and acorns, while also offering a lucrative cash crop in the form of bark that is utilized in the production of traditional corks (Acha & Newing, 2015).




Figure 5. The benefits of agroforestry.

According to Chenyang et al. (2021), perennial polycultures such as agroforestry exhibit more intrinsic stability when confronted with global market volatility and extreme climatic events, in comparison to annual monocultures. In the hypothetical scenario of a scarcity of fossil fuels, it is anticipated that mature fruit and nut trees will sustain their production with minimal disruption, albeit necessitating the substitution of labor for alternative inputs. Agroforests possess the capacity to store greenhouse gases, which are recognized as the primary drivers of global climate change (Wilson & Lovell, 2016; Raihan et al., 2022b; Jubair et al., 2023; Raihan, 2024c). Additionally, these agroforests have a heightened ability to withstand and adapt to the anticipated impacts of climate change. According to Rajanna et al. (2023), enhanced root systems and increased capacity for infiltration and water retention contribute to mitigating the effects of drought. Additionally, trees exhibit superior resilience to floods compared to field crops due to their capacity to extract surplus water from the soil and endure inundation. Despite being frequently disregarded, agroforestry offers supplementary cultural advantages. The preservation of nature is highly esteemed by several landowners due to its aesthetic appeal and perceived advantages, such as enhanced well-being and the tranquility associated with rural living (Tindale et al., 2023). According to a study conducted by Jiang et al. (2023), rural dwellers exhibit a preference for aesthetics that are enhanced by the use of measures such as vegetative buffers. Etongo et al. (2023) highlight several recreational activities available, such as bird watching, outdoor excursions, and hunting. The benefits of agroforestry are illustrated in Figure 5.

Agroforestry Adoption Challenges and Future Directions to Overcome

The prospects for agroforestry are promising, albeit not devoid of obstacles. The adoption of agroforestry has exhibited a notable deficiency, despite the existence of extensively demonstrated advantages (Syano et al., 2022). Various obstacles have been identified, such as the financial burden associated with setting up tree plantations, landowners' limited familiarity with tree cultivation, and the considerable time and expertise necessary for effective management (Irwin et al., 2023). According to Wienhold and Goulao (2023), extension employees and agricultural product merchants are commonly relied upon by farmers for acquiring knowledge about novel agricultural methods. However, it is worth noting that these experts generally lack formal training or practical expertise in the field of agroforestry. Furthermore, the absence of well-defined demonstration plots poses a challenge for landowners in observing the practical implementation of these systems (Zang et al., 2022). Given the intangible or long-term nature of numerous beneficial outcomes associated with agroforestry, landowners may encounter challenges in visualizing them (Jacobs et al., 2023).

The logistics associated with the harvest of edible goods, such as fruits and nuts, in agroforestry systems can pose significant challenges. In order to enhance the economic competitiveness of agroforestry systems, the implementation of mechanization may be necessary for larger-scale plantings (Korneeva & Belyaev, 2022). The complexity of the situation arises when numerous types of fruit or nut are cultivated simultaneously. According to Irwin et al. (2023), non-traditional markets and delayed rewards can also serve as deterrents. Previous studies have demonstrated that certain agroforestry systems, like silvopasture, exhibit economic viability and generate profits. However, other practices such as biomass plantings or riparian buffers may require the establishment of markets that provide compensation for the ecosystem services, they offer in order to be financially feasible (Mosquera-Losada et al., 2023; Ntawuruhunga et al., 2023). Additionally, the process of social change and networking will also be influential as attitudes shift to embrace alternative approaches (Annosi et al., 2022).

In light of the aforementioned obstacles, several techniques have been presented with the aim of advancing the field of agroforestry. Potential policy adjustments may involve the augmentation of financial resources allocated towards government cost-sharing initiatives aimed at facilitating the implementation of sustainable practices. Additionally, the provision of incentives, such as credits, for the provision of environmental services, including

but not limited to pollination and carbon sequestration, might be considered. The existing programs offered by the United States Department of Agriculture (USDA) through the Natural Resources Conservation Service (NRCS) and the Farm Service Agency (FSA) frequently include provisions that prohibit the harvesting of land designated for conservation purposes. However, the implementation of agroforestry systems has the potential to allow for the cultivation of harvestable products while still maintaining the conservation objectives. The implementation of a policy modification that permits the non-destructive extraction of consumable products from agroforestry systems could potentially incentivize a greater number of farmers to embrace agroforestry methods, hence resulting in improved conservation outcomes. While it is justifiable that a significant portion of government financing is allocated to support prominent cropping systems like maize and soybean, it is worth considering that agroforestry possesses the potential to mitigate the adverse impacts associated with these systems. Consequently, it is advisable to allocate greater attention and resources to agroforestry practices. A portion of this assistance has the potential to be allocated towards educational initiatives, specifically through the implementation of extension and university programs. Indeed, education emerges as a critical determinant for the adoption of conservation techniques, since numerous research investigating the adoption phenomenon consistently identify limited access to knowledge and technical support as a prominent obstacle.

The potential for expanding the output capacity of agroforestry systems remains largely untapped. Further investigation is required to examine the utilization of trees and shrubs in the production of commercially viable goods. In recent times, there has been a surge in interest in the advancement of multifunctional, consumable polycultures that emulate natural ecosystems, such as the indigenous oak savannas found in the Midwest region. These polycultures encompass the cultivation of various crops in a stacked arrangement, enabling the utilization of diverse ecological niches and the generation of multiple revenue streams. Field trials were conducted at the University of Illinois at Champaign-Urbana to investigate the cultivation of a combination of chestnuts, hazelnuts, apples, currants, and raspberries. The inclusion of control plots in a conventionally managed corn and soy rotation enables the opportunity to conduct a comprehensive examination of various environmental, ecological, and economic variables for comparison purposes. A comprehensive and repeatable investigation was conducted to examine various spatial arrangements of polycultures in comparison to monocultures of individual species, as they would typically be cultivated in a commercial orchard setting. Additionally, the study allowed for a comparison between these polycultures and a corn/soybean rotation. The treatments encompass the incorporation of indigenous trees and shrubs that possess consumable produce, such as aronia, elderberry, pecan, pawpaw, persimmon, plum, and serviceberry. This study investigates the potential scope of cultivating native culinary plants within conservation easements that require the exclusive utilization of indigenous species.

Conclusion

This paper aims to provide a review of agroforestry toward sustainable and resilient agriculture for the future world. Numerous strategies have been suggested to effectively and durably address the challenge of providing nourishment to an expanding global populace. Organic farming exhibits the potential to reduce the reliance on agrichemicals and enhance specific environmental and human health indicators. Conversely, advocates of conventional farming systems highlight the benefits associated with the utilization of genetic engineering, fertilizers, and pest control methods to enhance crop productivity. The implementation of broader methods encompasses various approaches, such as the restriction of farmland expansion through the prevention of deforestation, the reduction of food waste, the adoption of a less meat-intensive diet, the closure of yield gaps in underperforming cropland within emerging nations, and the enhancement of resource efficiency pertaining to water, fertilizer, and fuel utilization. These aforementioned endeavors, together with additional strategies, will be

imperative components of a comprehensive approach in order to effectively and enduringly address global food security.

The natural environment yields its abundance without necessitating the use of plowing, fertilizers, or pest control measures, hence obviating the need for any external inputs. The system operates only on solar energy and does not produce any detrimental waste byproducts. The presence of biological diversity enables the capacity for dynamic adaptation in response to environmental changes. By emulating the operation of natural ecosystems, agricultural systems have the potential to enhance their stability and resilience. Constructing such a system undoubtedly presents a formidable undertaking, necessitating a diverse array of technologies. Agroforestry has the potential to serve as a progressive approach in the realm of sustainable agriculture. It achieves this by advocating for and implementing integrated and biodiverse practices that aim to enhance crop yields, mitigate adverse impacts, and deepen our comprehension of the intricate interdependencies inherent in augmenting food production while mitigating harm. Agroforestry could lead to long-term sustainable change with a balance between short-term economic benefits and long-term sustainability goals. Sustainable agroforestry systems have the potential to help farmers harness the interactions occurring between the different components of the system for a multitude of benefits such as increased yield, environmental benefits, and animal welfare. Therefore, agroforestry should be given higher priority as a nature-based solution in policies and programs aimed at ecosystem restoration, land degradation neutrality, and climate change mitigation goals, particularly for developing countries.

Declaration

Acknowledgment: N/A

Funding: This research received no funding.

Conflict of interest: The author declares no conflict of interest.

Authors contribution: Asif Raihan contributed to conceptualization, visualization, methodology, reviewing literature, extracting information, synthesizing, and manuscript writing.

Data availability: The author confirms that the data supporting the findings of this study are available within the article.

References

- Acha, A., & Newing, H. S. (2015). Cork oak landscapes, promised or compromised lands? A case study of a traditional cultural landscape in Southern Spain. *Human Ecology*, 43, 601-611.
- Ahmed, Z., & Ambinakudige, S. (2023). Does land use change, waterlogging, and salinity impact on sustainability of agriculture and food security? Evidence from southwestern coastal region of Bangladesh. *Environmental Monitoring and Assessment*, 195(1), 74.
- Annosi, M. C., Ráez, R. M. O., Appio, F. P., & Del Giudice, T. (2022). An integrative review of innovations in the agricultural sector: The roles of agency, structure, and their dynamic interplay. *Technological Forecasting and Social Change*, 185, 122035.
- Arnhold, S., Lindner, S., Lee, B., Martin, E., Kettering, J., Nguyen, T. T., ... & Huwe, B. (2014). Conventional and organic farming: Soil erosion and conservation potential for row crop cultivation. *Geoderma*, 219, 89-105.
- Aryal, K., Maraseni, T., & Apan, A. (2023). Transforming agroforestry in contested landscapes: A win-win solution to trade-offs in ecosystem services in Nepal. *Science of The Total Environment*, 857, 159301.

- Balasundram, S. K., Shamshiri, R. R., Sridhara, S., & Rizan, N. (2023). The Role of Digital Agriculture in Mitigating Climate Change and Ensuring Food Security: An Overview. *Sustainability*, 15(6), 5325.
- Beaumelle, L., Giffard, B., Tolle, P., Winter, S., Entling, M. H., Benítez, E., ... & Rusch, A. (2023). Biodiversity conservation, ecosystem services and organic viticulture: A glass half-full. *Agriculture, Ecosystems & Environment*, 351, 108474.
- Bedeke, S. B. (2023). Climate change vulnerability and adaptation of crop producers in sub-Saharan Africa: A review on concepts, approaches and methods. *Environment, Development and Sustainability*, 25(2), 1017-1051.
- Begum, R. A., Raihan, A., & Said, M. N. M. (2020). Dynamic impacts of economic growth and forested area on carbon dioxide emissions in Malaysia. *Sustainability*, 12(22), 9375.
- Behera, B., Das, T. K., Raj, R., Ghosh, S., Raza, M. B., & Sen, S. (2021). Microbial consortia for sustaining productivity of non-legume crops: prospects and challenges. *Agricultural Research*, 10, 1-14.
- Benita, F. (2021). Human mobility behavior in COVID-19: A systematic literature review and bibliometric analysis. *Sustainable Cities and Society*, 70, 102916.
- Bishaw, B., Soolanayakanahally, R., Karki, U., & Hagan, E. (2022). Agroforestry for sustainable production and resilient landscapes. *Agroforestry Systems*, 96(3), 447-451.
- Brick, E. S. R., Holland, J., Anagnostou, D. E., Brown, K., & Desmulliez, M. P. (2022). A review of agroforestry, precision agriculture, and precision livestock farming—The case for a data-driven agroforestry strategy. *Frontiers in Sensors*, 3, 998928.
- Caporali, F. (2021). Sustainable Agriculture Through Ecological Intensification. *Ethics and Sustainable Agriculture: Bridging the Ecological Gaps*, 123-258.
- Chamberlain, J., Small, C., & Baumflek, M. (2019). Sustainable forest management for nontimber products. *Sustainability*, 11(9), 2670.
- Chenyang, L., Currie, A., Darrin, H., & Rosenberg, N. (2021). Farming with Trees; Reforming US Farm Policy to Expand Agroforestry and Mitigate Climate Change. *Ecology LQ*, 48, 1.
- Chinthalapudi, D. P., Pokhrel, S., Kingery, W. L., Shankle, M. W., & Ganapathi Shanmugam, S. (2023). Exploring the Synergistic Impacts of Cover Crops and Fertilization on Soil Microbial Metabolic Diversity in Dryland Soybean Production Systems Using Biolog EcoPlates. *Applied Biosciences*, 2(3), 328-346.
- Crystal-Ornelas, R., Thapa, R., & Tully, K. L. (2021). Soil organic carbon is affected by organic amendments, conservation tillage, and cover cropping in organic farming systems: A meta-analysis. *Agriculture, Ecosystems & Environment*, 312, 107356.
- Dasgupta, R., Dhyani, S., Basu, M., Kadaverugu, R., Hashimoto, S., Kumar, P., ... & Mitra, P. (2023). Exploring indigenous and local knowledge and practices (ILKPs) in traditional jhum cultivation for localizing sustainable development goals (SDGs): a case study from Zunheboto district of Nagaland, India. *Environmental Management*, 72(1), 147-159.
- Davis, H., Magistrali, A., Butler, G., & Stergiadis, S. (2022). Nutritional benefits from fatty acids in organic and grass-fed beef. *Foods*, 11(5), 646.
- Derouiche, A., Achour, A., & Driss, M. R. (2023). Organochlorine pesticides and polychlorinated biphenyls in raw bovine milk from various dairy farms in Beja, Tunisia: contamination status, dietary intake, and health risk assessment for the consumers. *Environmental Science and Pollution Research*, 30(24), 65427-65439.
- Dipu, M. A., Jones, N. A., & Aziz, A. A. (2022). Drivers and barriers to uptake of regenerative agriculture in southeast Queensland: a mental model study. *Agroecology and Sustainable Food Systems*, 46(10), 1502-1526.
- Donley, N. (2019). The USA lags behind other agricultural nations in banning harmful pesticides. *Environmental Health*, 18(1), 1-12.

- Dupraz, C., Talbot, G., Marrou, H., Wery, J., Roux, S., Liagre, F., ... & Nogier, A. (2011). To mix or not to mix: evidences for the unexpected high productivity of new complex agrivoltaic and agroforestry systems. In *Proceedings of the 5th world congress of conservation agriculture: Resilient food systems for a changing world*.
- Englund, O., Börjesson, P., Mola-Yudego, B., Berndes, G., Dimitriou, I., Cederberg, C., & Scarlat, N. (2021). Strategic deployment of riparian buffers and windbreaks in Europe can co-deliver biomass and environmental benefits. *Communications Earth & Environment*, 2(1), 176.
- Etongo, D., Lafleur, H., & Vel, T. (2023). Community perceptions and attitudes towards the management of protected areas in Seychelles with Morne Seychellois National Park as case study. *World Development Sustainability*, 3, 100091.
- Fahad, S., Chavan, S. B., Chichaghare, A. R., Uthappa, A. R., Kumar, M., Kakade, V., ... & Poczai, P. (2022). Agroforestry systems for soil health improvement and maintenance. *Sustainability*, 14(22), 14877.
- Feng, L., Chi, B., & Dong, H. (2022). Cotton cultivation technology with Chinese characteristics has driven the 70-year development of cotton production in China. *Journal of Integrative Agriculture*, 21(3), 597-609.
- Francaviglia, R., Almagro, M., & Vicente-Vicente, J. L. (2023). Conservation agriculture and soil organic carbon: principles, processes, practices and policy options. *Soil Systems*, 7(1), 17.
- Frey, G. E., Chamberlain, J. L., & Jacobson, M. G. (2023). Producers, production, marketing, and sales of non-timber forest products in the United States: a review and synthesis. *Agroforestry Systems*, 97(3), 355-368.
- Gagliardi, L., Fontanelli, M., Frascioni, C., Sportelli, M., Antichi, D., Tramacere, L. G., ... & Raffaelli, M. (2022). Assessment of a Chain Mower Performance for Weed Control under Tree Rows in an Alley Cropping Farming System. *Agronomy*, 12(11), 2785.
- Gerhardt, C., Bröring, S., Strecker, O., Wustmans, M., Moretti, D., Breunig, P., ... & Förster, B. (2022). Framework for the Digital Transformation of the Agricultural Ecosystem. In *Handbook Digital Farming: Digital Transformation for Sustainable Agriculture* (pp. 59-108). Berlin, Heidelberg: Springer Berlin Heidelberg.
- Ghosh, D., John, E. A., & Wilkinson, A. (2023). Clever pest control? The role of cognition in biological pest regulation. *Animal Cognition*, 26(1), 189-197.
- Gou, Z., Yin, W., Asibi, A. E., Fan, Z., Chai, Q., & Cao, W. (2022). Improving the sustainability of cropping systems via diversified planting in arid irrigation areas. *Agronomy for Sustainable Development*, 42(5), 88.
- Graves, A. R., Burgess, P. J., Palma, J. H. N., Herzog, F., Moreno, G., Bertomeu, M., ... & Van den Briel, J. P. (2007). Development and application of bio-economic modelling to compare silvoarable, arable, and forestry systems in three European countries. *Ecological Engineering*, 29(4), 434-449.
- Gravesen, M. L., & Funder, M. (2022). *The Great Green Wall: An overview and lessons learnt* (No. 2022: 02). DIIS Working Paper.
- Haile, S., Palmer, M., & Otey, A. (2016). Potential of loblolly pine: switchgrass alley cropping for provision of biofuel feedstock. *Agroforestry Systems*, 90, 763-771.
- Hidalgo-Galvez, M. D., Barkaoui, K., Volaire, F., Matías, L., Cambrollé, J., Fernández-Rebollo, P., ... & Pérez-Ramos, I. M. (2022). Can trees buffer the impact of climate change on pasture production and digestibility of Mediterranean dehesas?. *Science of the Total Environment*, 835, 155535.
- Huang, I. Y., James, K., Thamthanakoon, N., Pinitjitsamut, P., Rattanamane, N., Pinitjitsamut, M., ... & Lowenberg-DeBoer, J. (2023). Economic outcomes of rubber-based agroforestry systems: a systematic review and narrative synthesis. *Agroforestry Systems*, 97(3), 335-354.
- Irwin, R., Short, I., Mohammadrezaei, M., & Dhuháin, Á. N. (2023). Increasing tree cover on Irish dairy and drystock farms: The main attitudes, influential bodies and barriers that affect agroforestry uptake. *Environmental Science & Policy*, 146, 76-89.

- Jacobs, H., Gupta, A., & Möller, I. (2023). Governing-by-aspiration? Assessing the nature and implications of including negative emission technologies (NETs) in country long-term climate strategies. *Global Environmental Change*, 81, 102691.
- Jacquet, F., Jeuffroy, M. H., Jouan, J., Le Cadre, E., Litrico, I., Malausa, T., ... & Huyghe, C. (2022). Pesticide-free agriculture as a new paradigm for research. *Agronomy for Sustainable Development*, 42(1), 8.
- Jayathilake, H. M., Prescott, G. W., Carrasco, L. R., Rao, M., & Symes, W. S. (2021). Drivers of deforestation and degradation for 28 tropical conservation landscapes. *Ambio*, 50(1), 215-228.
- Jiang, S., Ma, H., Yang, L., & Luo, S. (2023). The Influence of Perceived Physical and Aesthetic Quality of Rural Settlements on Tourists' Preferences—A Case Study of Zhaoxing Dong Village. *Land*, 12(8), 1542.
- Jinger, D., Kumar, R., Kakade, V., Dinesh, D., Singh, G., Pande, V. C., ... & Singhal, V. (2022). Agroforestry for controlling soil erosion and enhancing system productivity in ravine lands of Western India under climate change scenario. *Environmental Monitoring and Assessment*, 194(4), 267.
- Jones, M. W., Peters, G. P., Gasser, T., Andrew, R. M., Schwingshackl, C., Gütschow, J., ... & Le Quéré, C. (2023). National contributions to climate change due to historical emissions of carbon dioxide, methane, and nitrous oxide since 1850. *Scientific Data*, 10(1), 155.
- Jubair, A. N. M., Rahman, M. S., Sarmin, I. J., & Raihan, A. (2023). Tree diversity and regeneration dynamics toward forest conservation and environmental sustainability: A case study from Nawabganj Sal Forest, Bangladesh. *Journal of Agriculture Sustainability and Environment*, 2(2), 1-22.
- Kalogiannidis, S., Chatzitheodoridis, F., Kontsas, S., & Syndoukas, D. (2023). Impact of Bioenergy on Economic Growth and Development: An European Perspective. *International Journal of Energy Economics and Policy*, 13(3), 494.
- Khatri, P., Kumar, P., Shakya, K. S., Kirlas, M. C., & Tiwari, K. K. (2023). Understanding the intertwined nature of rising multiple risks in modern agriculture and food system. *Environment, Development and Sustainability*, 1-44.
- Kim, D. G., Kirschbaum, M. U., & Beedy, T. L. (2016). Carbon sequestration and net emissions of CH₄ and N₂O under agroforestry: Synthesizing available data and suggestions for future studies. *Agriculture, Ecosystems & Environment*, 226, 65-78.
- Korneeva, E. A., & Belyaev, A. I. (2022). Assessment of ecological and economic efficiency of agroforestry systems in arid conditions of the lower Volga. *Forests*, 13(8), 1248.
- Kumar, P., Uthappa, A. R., Chavan, S. B., Chichaghare, A. R., Debta, H., Bhat, S., & Dagar, J. C. (2023). Achieving Biodiversity Conservation, Livelihood Security and Sustainable Development Goals Through Agroforestry in Coastal and Island Regions of India and Southeast Asia. In *Agroforestry for Sustainable Intensification of Agriculture in Asia and Africa* (pp. 429-486). Singapore: Springer Nature Singapore.
- La, N., Bergkvist, G., Dahlin, A. S., Mulia, R., & Öborn, I. (2023). Agroforestry with contour planting of grass contributes to terrace formation and conservation of soil and nutrients on sloping land. *Agriculture, Ecosystems & Environment*, 345, 108323.
- Lamichhane, J. R., Alletto, L., Cong, W. F., Dayoub, E., Maury, P., Plaza-Bonilla, D., ... & Debaeke, P. (2023). Relay cropping for sustainable intensification of agriculture across temperate regions: Crop management challenges and future research priorities. *Field Crops Research*, 291, 108795.
- Li, T. (2021). Protecting the breadbasket with trees? The effect of the great plains shelterbelt project on agriculture. *Land Economics*, 97(2), 321-344.
- Lin, B. J., Li, R. C., Liu, K. C., Pelumi Oladele, O., Xu, Z. Y., Lal, R., ... & Zhang, H. L. (2023). Management-induced changes in soil organic carbon and related crop yield dynamics in China's cropland. *Global Change Biology*, 29(13), 3575-3590.

- Lind, L., Hasselquist, E. M., & Laudon, H. (2019). Towards ecologically functional riparian zones: A meta-analysis to develop guidelines for protecting ecosystem functions and biodiversity in agricultural landscapes. *Journal of environmental management*, 249, 109391.
- Liu, J., & Lobb, D. A. (2021). An overview of crop and crop residue management impacts on crop water use and runoff in the Canadian prairies. *Water*, 13(20), 2929.
- Lugo-Pérez, J., Hajian-Forooshani, Z., Perfecto, I., & Vandermeer, J. (2023). The importance of shade trees in promoting carbon storage in the coffee agroforest systems. *Agriculture, Ecosystems & Environment*, 355, 108594.
- Majeed, Y., Khan, M. U., Waseem, M., Zahid, U., Mahmood, F., Majeed, F., ... & Raza, A. (2023). Renewable energy as an alternative source for energy management in agriculture. *Energy Reports*, 10, 344-359.
- Mallareddy, M., Thirumalaikumar, R., Balasubramanian, P., Naseeruddin, R., Nithya, N., Mariadoss, A., ... & Vijayakumar, S. (2023). Maximizing water use efficiency in rice farming: A comprehensive review of innovative irrigation management technologies. *Water*, 15(10), 1802.
- Mancho, C., Diez-Pascual, S., Alonso, J., Gil-Díaz, M., & Lobo, M. C. (2023). Assessment of recovered struvite as a safe and sustainable phosphorous fertilizer. *Environments*, 10(2), 22.
- Martín-Lammerding, D., Gabriel, J. L., Zambrana, E., Santín-Montanyá, I., & Tenorio, J. L. (2021). Organic amendment vs. Mineral fertilization under minimum tillage: Changes in soil nutrients, soil organic matter, biological properties and yield after 10 years. *Agriculture*, 11(8), 700.
- Mie, A., Andersen, H. R., Gunnarsson, S., Kahl, J., Kesse-Guyot, E., Rembialkowska, E., ... & Grandjean, P. (2017). Human health implications of organic food and organic agriculture: a comprehensive review. *Environmental Health*, 16(1), 1-22.
- Miñarro, M., García, D., & Rosa-García, R. (2023). Pollination of exotic fruit crops depends more on extant pollinators and landscape structure than on local management of domestic bees. *Agriculture, Ecosystems & Environment*, 347, 108387.
- Monteiro, A., & Santos, S. (2022). Sustainable approach to weed management: The role of precision weed management. *Agronomy*, 12(1), 118.
- Montgomery, D. R., Biklé, A., Archuleta, R., Brown, P., & Jordan, J. (2022). Soil health and nutrient density: preliminary comparison of regenerative and conventional farming. *PeerJ*, 10, e12848.
- Mosquera-Losada, M. R., Santos, M. G. S., Gonçalves, B., Ferreiro-Domínguez, N., Castro, M., Rigueiro-Rodríguez, A., ... & Santiago-Freijanes, J. J. (2023). Policy challenges for agroforestry implementation in Europe. *Frontiers in Forests and Global Change*, 6, 1127601.
- Mousavi, M., Taki, M., Raeini, M. G., & Soheilifard, F. (2023). Evaluation of energy consumption and environmental impacts of strawberry production in different greenhouse structures using life cycle assessment (LCA) approach. *Energy*, 280, 128087.
- Mume, I. D., & Workalemahu, S. (2021). Review on windbreaks agroforestry as a climate smart agriculture practices. *American Journal of Agriculture and Forestry*, 9(6), 342-347.
- Nair, P. R., Kumar, B. M., Nair, V. D., Nair, P. R., Kumar, B. M., & Nair, V. D. (2021). Agroforestry Systems in The Temperate Zone. *An Introduction to Agroforestry: Four Decades of Scientific Developments*, 195-232.
- Nosratti, I., Korres, N. E., & Cordeau, S. (2023). Knowledge of Cover Crop Seed Traits and Treatments to Enhance Weed Suppression: A Narrative Review. *Agronomy*, 13(7), 1683.
- Ntawuruhunga, D., Ngowi, E. E., Mangi, H. O., Salanga, R. J., & Shikuku, K. M. (2023). Climate-smart agroforestry systems and practices: A systematic review of what works, what doesn't work, and why. *Forest Policy and Economics*, 150, 102937.

- Pantović, J. G., & Sečanski, M. (2023). Weed Control in Organic Farming. *Contemporary Agriculture*, 72(1-2), 43-56.
- Paul, C., Bartkowski, B., Dönmez, C., Don, A., Mayer, S., Steffens, M., ... & Helming, K. (2023). Carbon farming: Are soil carbon certificates a suitable tool for climate change mitigation?. *Journal of Environmental Management*, 330, 117142.
- Pearsons, K. A., Omondi, E. C., Zinati, G., Smith, A., & Rui, Y. (2023). A tale of two systems: Does reducing tillage affect soil health differently in long-term, side-by-side conventional and organic agricultural systems?. *Soil and Tillage Research*, 226, 105562.
- Pergner, I., & Lippert, C. (2023). On the effects that motivate pesticide use in perspective of designing a cropping system without pesticides but with mineral fertilizer—a review. *Agronomy for Sustainable Development*, 43(2), 24.
- Piato, K., Subía, C., Pico, J., Calderón, D., Norgrove, L., & Lefort, F. (2021). Organic farming practices and shade trees reduce pest infestations in robusta coffee systems in Amazonia. *Life*, 11(5), 413.
- Pimentel, D., & Burgess, M. (2014). An environmental, energetic and economic comparison of organic and conventional farming systems. *Integrated Pest Management: Pesticide Problems*, Vol. 3, 141-166.
- Poudel, S., Fike, J. H., & Pent, G. J. (2022). Hair cortisol as a measure of chronic stress in ewes grazing either hardwood silvopastures or open pastures. *Agronomy*, 12(7), 1566.
- Prairie, A. M., King, A. E., & Cotrufo, M. F. (2023). Restoring particulate and mineral-associated organic carbon through regenerative agriculture. *Proceedings of the National Academy of Sciences*, 120(21), e2217481120.
- Prout, J. M., Shepherd, K. D., McGrath, S. P., Kirk, G. J., & Haefele, S. M. (2021). What is a good level of soil organic matter? An index based on organic carbon to clay ratio. *European Journal of Soil Science*, 72(6), 2493-2503.
- Raihan, A. (2023a). Toward sustainable and green development in Chile: dynamic influences of carbon emission reduction variables. *Innovation and Green Development*, 2, 100038.
- Raihan, A. (2023b). A review of the global climate change impacts, adaptation strategies, and mitigation options in the socio-economic and environmental sectors. *Journal of Environmental Science and Economics*, 2(3), 36-58.
- Raihan, A. (2023c). Exploring Environmental Kuznets Curve and Pollution Haven Hypothesis in Bangladesh: The Impact of Foreign Direct Investment. *Journal of Environmental Science and Economics*, 2(1), 25-36.
- Raihan, A. (2023d). The dynamic nexus between economic growth, renewable energy use, urbanization, industrialization, tourism, agricultural productivity, forest area, and carbon dioxide emissions in the Philippines. *Energy Nexus*, 9, 100180.
- Raihan, A. (2023e). A review on the integrative approach for economic valuation of forest ecosystem services. *Journal of Environmental Science and Economics*, 2(3), 1-18.
- Raihan, A. (2023f). A review of tropical blue carbon ecosystems for climate change mitigation. *Journal of Environmental Science and Economics*, 2(4), 14-36.
- Raihan, A. (2023g). Sustainable development in Europe: A review of the forestry sector's social, environmental, and economic dynamics. *Global Sustainability Research*, 2(3), 72-92.
- Raihan, A. (2023h). A comprehensive review of artificial intelligence and machine learning applications in energy consumption and production. *Journal of Technology Innovations and Energy*, 2(4), 1–26.
- Raihan, A. (2023i). A concise review of technologies for converting forest biomass to bioenergy. *Journal of Technology Innovations and Energy*, 2(3), 10-36.
- Raihan, A. (2023j). An overview of the energy segment of Indonesia: present situation, prospects, and forthcoming advancements in renewable energy technology. *Journal of Technology Innovations and Energy*, 2(3), 37-63.

- Raihan, A. (2024a). A Systematic Review of Geographic Information Systems (GIS) in Agriculture for Evidence-Based Decision Making and Sustainability. *Global Sustainability Research*, 3(1), 1-24.
- Raihan, A. (2024b). Influences of foreign direct investment and carbon emission on economic growth in Vietnam. *Journal of Environmental Science and Economics*, 3(1), 1-17.
- Raihan, A. (2024c). The potential of agroforestry in South Asian countries towards achieving the climate goals. *Asian Journal of Forestry* 8(1), 1-17.
- Raihan, A., & Bijoy, T. R. (2023). A review of the industrial use and global sustainability of Cannabis sativa. *Global Sustainability Research*, 2(4), 1-29.
- Raihan, A., & Himu, H. A. (2023). Global impact of COVID-19 on the sustainability of livestock production. *Global Sustainability Research*, 2(2), 1-11.
- Raihan, A., Muhtasim, D. A., Farhana, S., Hasan, M. A. U., Paul, A., & Faruk, O. (2022a). Toward environmental sustainability: Nexus between tourism, economic growth, energy use and carbon emissions in Singapore. *Global Sustainability Research*, 1(2), 53-65.
- Raihan, A., Muhtasim, D. A., Farhana, S., Pavel, M. I., Faruk, O., & Mahmood, A. (2022b). Nexus between carbon emissions, economic growth, renewable energy use, urbanization, industrialization, technological innovation, and forest area towards achieving environmental sustainability in Bangladesh. *Energy and Climate Change*, 3, 100080.
- Raihan, A., Muhtasim, D. A., Farhana, S., Rahman, M., Hasan, M. A. U., Paul, A., & Faruk, O. (2023a). Dynamic linkages between environmental factors and carbon emissions in Thailand. *Environmental Processes*, 10, 5.
- Raihan, A., Pavel, M. I., Muhtasim, D. A., Farhana, S., Faruk, O., & Paul, A. (2023b). The role of renewable energy use, technological innovation, and forest cover toward green development: Evidence from Indonesia. *Innovation and Green Development*, 2(1), 100035.
- Raihan, A., & Tuspekova, A. (2022a). The nexus between economic growth, renewable energy use, agricultural land expansion, and carbon emissions: new insights from Peru. *Energy Nexus*, 6, 100067.
- Raihan, A., & Tuspekova, A. (2022b). Dynamic impacts of economic growth, renewable energy use, urbanization, industrialization, tourism, agriculture, and forests on carbon emissions in Turkey. *Carbon Research*, 1(1), 20.
- Raihan, A., & Tuspekova, A. (2022c). Dynamic impacts of economic growth, energy use, urbanization, tourism, agricultural value-added, and forested area on carbon dioxide emissions in Brazil. *Journal of Environmental Studies and Sciences*, 12(4), 794-814.
- Raihan, A., & Tuspekova, A. (2022d). Dynamic impacts of economic growth, energy use, urbanization, agricultural productivity, and forested area on carbon emissions: new insights from Kazakhstan. *World Development Sustainability*, 1, 100019.
- Raihan, A., & Tuspekova, A. (2022e). Nexus between energy use, industrialization, forest area, and carbon dioxide emissions: new insights from Russia. *Journal of Environmental Science and Economics*, 1(4), 1-11.
- Raihan, A., & Tuspekova, A. (2022f). Toward a sustainable environment: Nexus between economic growth, renewable energy use, forested area, and carbon emissions in Malaysia. *Resources, Conservation & Recycling Advances*, 15, 200096.
- Raihan, A., & Tuspekova, A. (2022g). Role of economic growth, renewable energy, and technological innovation to achieve environmental sustainability in Kazakhstan. *Current Research in Environmental Sustainability*, 4, 100165.
- Raihan, A., & Tuspekova, A. (2023a). The role of renewable energy and technological innovations toward achieving Iceland's goal of carbon neutrality by 2040. *Journal of Technology Innovations and Energy*, 2(1), 22-37.

- Raihan, A., & Tuspekova, A. (2023b). Towards net zero emissions by 2050: the role of renewable energy, technological innovations, and forests in New Zealand. *Journal of Environmental Science and Economics*, 2(1), 1-16.
- Rajanna, G. A., Suman, A., & Venkatesh, P. (2023). Mitigating Drought Stress Effects in Arid and Semi-Arid Agro-Ecosystems through Bioirrigation Strategies—A Review. *Sustainability*, 15(4), 3542.
- Read, C. (2022). *The Great Famine in Ireland and Britain's Financial Crisis* (Vol. 19). Boydell & Brewer.
- Ribas, M. P., García-Ulloa, M., Espunyes, J., & Cabezón, O. (2023). Improving the assessment of ecosystem and wildlife health: Microbiome as an early indicator. *Current Opinion in Biotechnology*, 81, 102923.
- Robertson, G. P., Paul, E. A., & Harwood, R. R. (2000). Greenhouse gases in intensive agriculture: contributions of individual gases to the radiative forcing of the atmosphere. *Science*, 289(5486), 1922-1925.
- Ruch, B., Hefner, M., & Sradnick, A. (2023). Excessive Nitrate Limits the Sustainability of Deep Compost Mulch in Organic Market Gardening. *Agriculture*, 13(5), 1080.
- Santoni, M., Verdi, L., Imran Pathan, S., Napoli, M., Dalla Marta, A., Dani, F. R., ... & Ceccherini, M. T. (2023). Soil microbiome biomass, activity, composition and CO₂ emissions in a long-term organic and conventional farming systems. *Soil Use and Management*, 39(1), 588-605.
- Scavo, A., Fontanazza, S., Restuccia, A., Pesce, G. R., Abbate, C., & Mauromicale, G. (2022). The role of cover crops in improving soil fertility and plant nutritional status in temperate climates. A review. *Agronomy for Sustainable Development*, 42(5), 93.
- Sereke, F., Graves, A. R., Dux, D., Palma, J. H., & Herzog, F. (2015). Innovative agroecosystem goods and services: key profitability drivers in Swiss agroforestry. *Agronomy for sustainable development*, 35, 759-770.
- Shah, S. A. H., Bilal, A., Ahmad, M. M., & Bukhari, S. S. (2022). Deforestation is causing a great loss in avian diversity in Pakistan. *American Journal of Zoology*, 5(3), 24-29.
- Sivaranjani, S., & Panwar, V. P. (2023). Carbon sequestration in agroforestry systems. In *Agricultural Soil Sustainability and Carbon Management* (pp. 207-227). Academic Press.
- Smith, M. M., Bentrup, G., Kellerman, T., MacFarland, K., Straight, R., Ameyaw, L., & Stein, S. (2022). Silvopasture in the USA: A systematic review of natural resource professional and producer-reported benefits, challenges, and management activities. *Agriculture, Ecosystems & Environment*, 326, 107818.
- Sollen-Norrin, M., Ghaley, B. B., & Rintoul, N. L. J. (2020). Agroforestry benefits and challenges for adoption in Europe and beyond. *Sustainability*, 12(17), 7001.
- Stokes, A., Bocquého, G., Carrere, P., Salazar, R. C., Deconchat, M., Garcia, L., ... & Thomas, M. (2023). Services provided by multifunctional agroecosystems: Questions, obstacles and solutions. *Ecological Engineering*, 191, 106949.
- Subbulakshmi, V., Sheetal, K. R., Noor Mohamed, M. B., Renjith, P. S., & Kala, S. (2023). Arid Agroforestry for Thar Desert. In *Natural Resource Management in the Thar Desert Region of Rajasthan* (pp. 155-192). Cham: Springer International Publishing.
- Syano, N. M., Nyangito, M. M., Wasonga, O. V., & Kironchi, G. (2022). Agroforestry practices and factors influencing their adoption by communities in the drylands of Eastern Kenya. *Agroforestry Systems*, 96(8), 1225-1235.
- Szczepanek, M., Siwik-Ziomek, A., Lemańczyk, G., Lamparski, R., & Graczyk, R. (2023). Effect of reduced tillage on soil enzyme activity, pests pressure and productivity of organically grown spring wheat species. *Agronomy*, 13(2), 287.
- Tawfik, G. M., Dila, K. A. S., Mohamed, M. Y. F., Tam, D. N. H., Kien, N. D., Ahmed, A. M., & Huy, N. T. (2019). A step by step guide for conducting a systematic review and meta-analysis with simulation data. *Tropical medicine and health*, 47(1), 1-9.

- Taylor, J. R., & Lovell, S. T. (2021). Designing multifunctional urban agroforestry with people in mind. *Urban Agriculture & Regional Food Systems*, 6(1), e20016.
- Tei, F., De Neve, S., de Haan, J., & Kristensen, H. L. (2020). Nitrogen management of vegetable crops. *Agricultural Water Management*, 240, 106316.
- Telwala, Y. (2023). Unlocking the potential of agroforestry as a nature-based solution for localizing sustainable development goals: A case study from a drought-prone region in rural India. *Nature-Based Solutions*, 3, 100045.
- Temegne, N. C., Ngome, A. F., Agendia, A. P., & Youmbi, E. (2021). Agroecology for agricultural soil management. *Sustainable intensification for agroecosystem services and management*, 267-321.
- Thiesen, T., Bhat, M. G., Liu, H., & Rovira, R. (2022). An ecosystem service approach to assessing agroecosystems in urban landscapes. *Land*, 11(4), 469.
- Tindale, S., Vicario-Modroño, V., Gallardo-Cobos, R., Hunter, E., Miškolci, S., Price, P. N., ... & Frewer, L. J. (2023). Citizen perceptions and values associated with ecosystem services from European grassland landscapes. *Land Use Policy*, 127, 106574.
- Udawatta, R. P., & Jose, S. (2011). Carbon sequestration potential of agroforestry practices in temperate North America. *Carbon sequestration potential of agroforestry systems: opportunities and challenges*, 17-42.
- Valenzuela, H. (2023). Ecological Management of the Nitrogen Cycle in Organic Farms. *Nitrogen*, 4(1), 58-84.
- Vassar, M., Atakpo, P., & Kash, M. J. (2016). Manual search approaches used by systematic reviewers in dermatology. *Journal of the Medical Library Association: JMLA*, 104(4), 302.
- Viñals, E., Maneja, R., Rufi-Salís, M., Martí, M., & Puy, N. (2023). Reviewing social-ecological resilience for agroforestry systems under climate change conditions. *Science of the total environment*, 869, 161763.
- Wienhold, K., & Goulao, L. F. (2023). The Embedded Agroecology of Coffee Agroforestry: A Contextualized Review of Smallholder Farmers' Adoption and Resistance. *Sustainability*, 15(8), 6827.
- Wijerathna-Yapa, A., & Pathirana, R. (2022). Sustainable Agro-Food Systems for Addressing Climate Change and Food Security. *Agriculture*, 12(10), 1554.
- Wilson, M. H., & Lovell, S. T. (2016). Agroforestry—The next step in sustainable and resilient agriculture. *Sustainability*, 8(6), 574.
- Xu, S., Wang, R., Gasser, T., Ciais, P., Peñuelas, J., Balkanski, Y., ... & Zhang, R. (2022). Delayed use of bioenergy crops might threaten climate and food security. *Nature*, 609(7926), 299-306.
- Yousefi, A., Ghahramany, L., Ghazanfari, H., Pulido, F., & Moreno, G. (2020). Biometric indices of wild pistachio (*Pistacia atlantica* Desf.) trees under resin extraction in Western Iran. *Agroforestry Systems*, 94, 1977-1988.
- Yuan, X., Wang, Y., Ji, P., Wu, P., Sheffield, J., & Otkin, J. A. (2023). A global transition to flash droughts under climate change. *Science*, 380(6641), 187-191.
- Zahoor, I., & Mushtaq, A. (2023). Water Pollution from Agricultural Activities: A Critical Global Review. *Int. J. Chem. Biochem. Sci*, 23, 164-176.
- Zang, L., Wang, Y., Ke, J., & Su, Y. (2022). What drives smallholders to utilize socialized agricultural services for farmland scale management? Insights from the perspective of collective action. *Land*, 11(6), 930.
- Zhang, S., Huang, G., Zhang, Y., Lv, X., Wan, K., Liang, J., ... & Hu, F. (2023). Sustained productivity and agronomic potential of perennial rice. *Nature Sustainability*, 6(1), 28-38.